

From the Department of Laboratory Medicine, Division of Pathology
Karolinska Institutet, Karolinska University Hospital, Huddinge
Stockholm, Sweden

**TARTRATE RESISTANT ACID PHOSPHATASE 5A: A POTENTIAL REGULATOR OF
ADIPOCYTE CELL NUMBER AND DIFFERENTIATION IN WHITE ADIPOSE TISSUE**

Christina Patlaka



**Karolinska
Institutet**

Stockholm 2015

Cover: Confocal microscopy image of pre-adipocytes treated with TRAP 5a at 4°C and stained for TRAP 5a (red)

All previously published papers were reproduced with permission from the publisher.

Published by Karolinska Institutet.

Printed by Eprint AB 2015

© Christina Patlaka, 2015

ISBN 978-91-7549-901-7

Tartrate resistant acid phosphatase 5a: A potential regulator of adipocyte cell number and differentiation in white adipose tissue

THESIS FOR DOCTORAL DEGREE (Ph.D.)

By

Christina Patlaka

Principal Supervisor:

Professor Göran Andersson
Karolinska Institutet
Department of Laboratory Medicine
Division of Pathology

Co-supervisor(s):

Pernilla Lång PhD
Karolinska Institutet
Department of Laboratory Medicine
Division of Pathology

Opponent:

Professor Peter Strålfors
Linköping University
Department of Clinical and Experimental
Medicine
Division of Cell Biology

Examination Board:

Professor Rachel Fisher
Karolinska Institutet
Department of Medicine
Division of Atherosclerosis Research Unit

Associate Professor Jurga Laurencikiene
Karolinska Institutet
Department of Medicine
Division of Lipid laboratory

Professor Anders Jacobsson
Stockholm University
Department of The Wenner-Gren Institute
Division of Physiology

To my family

ABSTRACT

Tartrate-resistant acid phosphatase (TRAP) exists in two isoforms, TRAP 5a which is monomeric and TRAP 5b which is a dimer generated by proteolytic cleavage of TRAP 5a, that exhibit different functions and localizations. TRAP 5a is expressed by adipose tissue macrophages and secreted into the extracellular environment and has been shown to lead to hyperplastic insulin-sensitive obesity when over-expressed in mice. In bone, TRAP is suggested to interact with the heparan sulfate proteoglycan (HSPG) glypican-4. In humans, TRAP 5a serum levels and TRAP in adipose tissue correlate to BMI. TRAP 5a has been shown to have a stimulatory effect on the proliferation and differentiation of pre-adipocytes, pre-osteoblasts and hematopoietic cells but the mechanism of action remains unknown. The aims of this thesis were (1) to develop an ELISA for quantification of human TRAP 5a and to evaluate TRAP 5a levels in lean vs. obese individuals and (2) to investigate the mechanism of action of TRAP 5a in pre-adipocytes and the effects on the homeostasis-related events such as cell cycle entry, differentiation and migration using the pre-adipocyte cell line 3T3-L1. A TRAP 5a ELISA was developed by generation of monoclonal antibodies specific for TRAP 5a and used to measure TRAP 5a in serum from lean and obese females. TRAP 5a in serum of obese females was reduced compared to lean but a positive correlation between BMI >30 and serum levels of TRAP 5a was identified. Regarding effect of TRAP 5a on 3T3-L1 cells, TRAP 5a was shown to bind to the components of extracellular matrix (ECM) heparan sulfate and nidogen-2. TRAP 5a was shown to bind to the pre-adipocyte cell membrane and to co-localize with HSPG glypican-4. After cell surface binding, TRAP 5a is endocytosed in a caveolae-mediated manner and co-localizes with early endosomal marker Rab-7. TRAP 5a was also observed in multivesicular bodies but did not co-localize with lysosomal marker LAMP-1. Mesenchymal stem cells, fibroblasts and pre-osteoblasts were also tested for their ability to bind TRAP 5a and only pre-osteoblasts exhibited TRAP 5a endocytosis. TRAP 5a was shown to enhance the number of cells entering S phase after cell cycle arrest by serum deprivation and to increase cyclin D1 levels. In a phosphorylation signaling array for proteins of the Akt pathway it was shown that TRAP 5a causes increased inactivation phosphorylation of GSK3 β that is most likely not dependent on Akt since Akt activation phosphorylation was reduced after TRAP 5a treatment. Proteins controlling cell cycle arrest and apoptosis p53, p38 and Bad exhibited altered phosphorylation pattern after TRAP 5a treatment that could potentially lead to their inactivation. TRAP 5a treatment induced changes in morphology of pre-adipocytes and enhanced their migration. Focal adhesion protein (FAK), paxillin and actin were down-regulated after TRAP treatment indicating cytoskeletal changes and changes in the adhesion pattern. TRAP 5a treatment lead to enhanced lipid droplet acquisition after 2 days of differentiation induction in pre-adipocytes. In summary, TRAP 5a interacts with components of the ECM, binds to the surface of pre-adipocytes and co-localizes with glypican-4 and is endocytosed in a caveolae-dependent manner. Treatment of pre-adipocytes with TRAP 5a leads to enhanced cell numbers, increased migration, morphological changes and increased differentiation, all consistent with a growth factor-like role for TRAP 5a in adipose tissue.

LIST OF SCIENTIFIC PAPERS

- I. **Christina Patlaka**, Staffan Paulie, Laia Mira Pascual, Peter Arner, Pernilla Lång, Göran Andersson. Development of a human tartrate resistant acid phosphatase 5a specific ELISA. Manuscript
- II. **Christina Patlaka**, Hong Anh Mai, Pernilla Lång, Göran Andersson. The growth factor-like adipokine tartrate -resistant acid phosphatase 5a interacts with the rod G3 domain of adipocyte - produced nidogen-2. *Biochem Biophys Res Commun.*, 2014, 454, 3, 446-452
- III. **Christina Patlaka**, Heike Becker, Maria Norgård, Staffan Paulie, Annica Nordvall-Bodell, Pernilla Lång, Göran Andersson. Caveolae-mediated endocytosis of the glucosaminoglycan- interacting adipokine tartrate resistant acid phosphatase 5a in adipocyte progenitor lineage cells. *Biochim Biophys Acta*. 2014 Mar;1843(3):495-507.
- IV. **Christina Patlaka**, Laia Mira Pascual, Pernilla Lång, Göran Andersson. Tartrate resistant acid phosphatase 5a regulates cell number and motility in 3T3-L1 pre-adipocytes. Manuscript

CONTENTS

1	Introduction	1
1.1	Tissue homeostasis	1
1.1.1	How is cell communication mediated?.....	1
1.1.2	Major endocytosis route for growth factors	2
1.1.3	Cell number regulation.....	3
1.1.4	Cell recruitment and migration	5
1.1.5	Cell differentiation	6
1.2	Adipose tissue homeostasis.....	7
1.2.1	Adipose tissue as an endocrine and metabolic organ.....	7
1.2.2	Mesenchymal stem cells	8
1.2.3	From pre-adipocytes to adipocytes	9
1.2.4	Macrophages of adipose tissue	11
1.2.5	Adipose tissue changes in obesity	12
1.3	Tartrate- resistant acid phosphatase	13
1.3.1	TRAP history and properties	13
1.3.2	TRAP expression and localization	14
1.3.3	TRAP in disease	15
1.3.4	Regulation of cell number and differentiation by TRAP.....	16
1.3.5	Regulation of migration by TRAP.....	16
1.3.6	TRAP 5a in adipose tissue	16
1.3.7	Working hypothesis	17
2	AIMS of the thesis.....	18
3	Remarks on methodology	19
3.1	TRAP 5a ELISA development.....	19
3.2	Pull down assays.....	19
3.3	ICC/ 2-step indirect staining	19
3.4	Image analysis	20
3.5	Cell cycle analysis	20
3.6	Phosphorylation signalling assays	21
4	Results and discussion.....	23
4.1	Paper I.....	23
4.2	Paper II.....	24
4.3	Paper III	25
4.4	Paper IV	27
4.5	General Conclusions and discussion.....	28
4.6	Revised working hypothesis	29
4.7	Future Perspectives and clinical significance	31
5	Acknowledgements	33
6	References	35

LIST OF ABBREVIATIONS

aP2	Adipocyte protein 2	IL-4	Interleukin- 4
APPL	Amyloid protein precursor-like	JNK	c-Jun N-terminal kinase
ATM	Adipose tissue macrophages	LAMP-1	Lysosomal-associated membrane protein 1
BMI	Body mass index	LRP	Low-density lipoprotein receptor-related protein
BMP	Bone morphogenetic protein	MCP-1	Monocyte chemoattractant protein-1
C/EBPα	CCAAT/enhancer binding protein α	MMP	Matrix metalloproteinase
cAMP	Cyclic adenosine monophosphate	MSC	Mesenchymal stem cells
CCL	Chemokine (C-C motif) ligand	mTOR	Mechanistic target of rapamycin
Cdk	Cyclin -dependent kinase	MVB	Multi-vesicular bodies
CHD7 protein 7	Chromodomain-helicase-DNA-binding protein 7	NLK	Nemo-like kinase
COP-I	Coat protein I	PIK3	Phosphoinositide 3-kinase
CXCL	Chemokine (C-X-C motif) ligand	PIP₂	Phosphatidylinositol 4,5-bisphosphate
ECM	Extracellular matrix	PPARγ α	Peroxisome proliferator-activated receptor α
EEA	Early endosomal antigen	Pref-1	Pre-adipocyte factor 1
EGF	Epidermal growth factor	RAP1	Ras-proximate-1
EPAC cAMP	Exchange protein directly activated by cAMP	Rb	Retinoblastoma
ER	Endoplasmic reticulum	RTK	Receptor tyrosine kinases
FAK	Focal adhesion kinase	SARA	SMAD Anchor for Receptor Activation
FFA	Free fatty acid	SDF-1	Stromal cell-derived factor 1
FGF	Fibroblast growth factor	SNARE	SNAP Receptor
GLUT-4	Glucose transporter type 4	Stat 3	Signal transducer and activator of transcription 3
GPCR	G-protein coupled receptor	TG	Triglycerides
GSK3β	Glycogen synthase kinase 3 β	TGF-β	Tumor growth factor - β
GTP	Guanosine-5'-triphosphate	TIMP3	Tissue inhibitor of metalloproteinases 3
HGF	Hepatocyte growth factor	TNF-α	Tumor necrosis factor- α
HSP27	Heat shock protein 27	VCAM	Vascular cell-adhesion molecule 1
HSPG	Heparan sulfate proteoglycan	Wnt	Wingless integration-1
ICAM	Intracellular adhesion molecular 1		
IGF	Insulin-like growth factor		

1 INTRODUCTION

1.1 TISSUE HOMEOSTASIS

Tissue homeostasis refers to the maintenance of tissue in a functional state and is regulated by a variety of biological events regulating the life span of cells such as cell proliferation i.e. cell division, differentiation i.e. cell maturation, migration i.e. cell movement and cell death.

Tissues are multicellular structures comprised of different cell types. Tissue-specific cells are differentiated cells that have a particular function that can vary depending on the type of tissue e.g. a metabolic function like e.g. hepatocytes in liver or storage function (adipocytes in adipose tissue) or structural/barrier function such as keratinocytes in skin. Apart from differentiated cells, tissues contain progenitor/precursor cells, committed to a certain cell lineage, but not fully differentiated and retaining the ability, although reduced compared to e.g. stem cells, to proliferate (e.g. pre-adipocytes in adipose tissue). A fraction of stem cells named adult stem cells reside in tissue and serve as a reservoir pool for tissue regeneration of dying cells and maintain homeostasis in case of damage. Immune cells are additionally an important part of a tissue. Apart from dendritic cells and mast cells, resident macrophages are present in the stroma and are responsible for clearing of cells undergoing apoptosis and mediating the inflammatory response during tissue damage.

1.1.1 How is cell communication mediated?

Tissue homeostasis and the events described above (cell number regulation, cell migration and differentiation) are dependent on soluble signals from the extracellular environment (ECM), interaction of the cells with the ECM as well as with other cells.

ECM is involved in tissue homeostasis and cellular events in various ways. Growth factors can bind to the ECM and remain there until ECM degradation leads to their release and binding to the cells. Growth factors bind by the use of heparan sulfate (HS) of the ECM e.g. epidermal growth factor (EGF) and hepatocyte growth factor (HGF) and on heparan sulfate proteoglycans (HSPGs). Degradation of the ECM can lead to release of these factors once they are needed. Other growth factors, e.g. fibroblast growth factor (FGF) and transforming growth factor (TGF) use the proteoglycans (PG) of the ECM for binding to the cells, thus ECM facilitates the binding ¹. In other cases, the binding of growth factors to ECM serves a different purpose that is to make the growth factors available to the cells. Moreover, proteins of the ECM such as laminin contain intrinsic sequences (e.g. EGF-like domains) and have been suggested to directly bind to cell receptors and induce cell signalling ¹.

Mesenchymal stem cell (MSC) commitment to a certain cell lineage is also dependent on factors of the ECM that will provide the appropriate signals for the cells to differentiate. Formation of adipocyte precursors (pre-adipocytes) is impaired by the presence of the ECM component fibronectin and the binding to its receptor integrin $\alpha5/\beta1$ ². This binding causes the levels of the Rho GTPase RAC to remain high, thus inhibiting adipocyte differentiation. Rho GTPases are important regulators of MSC commitment as they influence the shape of the cells and for e.g. adipocytes, changes in shape are crucial for differentiation ². The extracellular environment can also be altered by the activation of matrix metalloproteinases (MMPs) and tissue inhibitors of MMPs (TIMPs) and influence cell lineage commitment, e.g. MMP14 and TIMP3 are involved in adipocyte lineage commitment ².

1.1.2 Major endocytosis route for growth factors

Once bound to their receptors, growth factors can follow various routes to complete the downstream signalling. Endocytosis can lead to different outcomes, such as signal reduction by receptor degradation, downstream signalling by interaction with proteins of the endocytic vesicles or signalling enhancement by receptor recycling.

The main mechanism of growth factor endocytosis are the clathrin- dependent and clathrin - independent endocytosis (Figure 1). One of the most common mechanisms of clathrin-dependent endocytosis is via β -arrestins that bind to activated (phosphorylated due to ligand binding) receptors, e.g. G-protein coupled receptors (GPCRs) and clathrin and adaptor protein 2 (AP2) thus forming an endocytic complex ³.

The most common clathrin- independent endocytosis pathways is mediated via caveolae. Caveolin-1 interaction with cholesterol and phosphorylation of caveolin-1 are two of the mechanisms that are required for the formation of caveolae ^{4,5}. Both clathrin- and caveolae-mediated endocytosis require the presence of dynamin for the formation of clathrin- and caveolae-coated pits and for later invagination of vesicles by hydrolyzing guanosine-5'-triphosphate (GTP) ^{6,7}. Downstream endocytosis can follow different routes. The first step for both clathrin- and caveolae- mediated endocytosis is the formation of early endosomes, where signalling transduction can take place via components Rab5 together with early endosomal antigen (EEA) (canonical endosome signalling) or APPL by formation of complex with Protein kinase B (PKB) also known as Akt and Glycogen synthase kinase 3 (GSK3 β) leading to its phosphorylation and downstream signalling ⁸. After early endosomes, receptor signalling can follow alternative routes. Receptor can be transported back to the cell membrane via recycling endosomes that contain Rab-4 or Rab-11. Duration of signalling can be determined by the action of recycling endosomes ⁸. Apart from recycling, receptors can be translocated to late endosomes and multivesicular endosomes/bodies for further transfer to the lysosomes for degradation but also for further signal enhancement. Late endosomes are characterized by the presence of Rab-7 that replaces the early endosome component Rab-5 ⁹.

For many growth factors, endocytosis takes place in order to sustain the signal by receptor recycling or in other cases to remove the ligand from the cell surface and transfer it to the lysosomes for degradation and signal reduction/deactivation. The endocytosis route can also be dependent on the ligand concentration. For example, EGF can use clathrin- dependent endocytosis for signal transduction at low concentrations but caveolae-dependent endocytosis for signal down-regulation at high concentrations ¹⁰. For other types of signalling, e.g. wingless/integration-1 (wnt) signalling, endocytosis via clathrin can lead to signal down-regulation, while clathrin-independent endocytosis to sustainment of signalling ¹¹.

Endocytosis can serve other purposes such as transmitting the signal from the cell membrane to specialized compartments where signalling is required. This is of particular importance in processes such as motility and migration of cells, where it has been shown that clathrin- and Rab5-mediated endocytosis is responsible for localized signalling of Rac that leads to actin re-organization and cell migration ¹². Finally, endocytosis alone can be determining for signal transduction, e.g. TGF- β binds to its receptor thus leading to its activation by phosphorylation and formation of endosomes, it causes the recruitment of SMAD2 by SMAD Anchor for Receptor Activation (SARA). This enables the phosphorylation of SMAD2 by TGF β receptor (TGF β R) and its translocation to the nucleus for initiation of transcription ^{13,14}. Multivesicular bodies (MVB) are actively participating in and are vital for signalling. GSK3 β signalling via Wnt is dependent upon endocytosis and MVB formation ¹⁵.




Figure 1. Endocytosis routes. Endocytic vesicles containing clathrin or caveolin and lipid rafts. After endocytosis there is formation of early endosomes that lead to recycling endosomes and transfer again to the membrane or to multivesicular bodies and late endosomes leading to lysosomal degradation.

1.1.2.1 Growth factor secretion from cells that regulate signals

Upon translation most growth factors and proteins follow conventional secretory pathways mediated via endoplasmic reticulum (ER)/golgi. Newly synthesized proteins can be transferred to ER and exit in coat-protein II (COP-II) vesicles to enter golgi and exit in coat-protein I (COP-I) coated vesicles that in their turn use SNAP Receptor (SNARE) complexes and a series of fusion of vesicles to exit through the cell membrane¹⁶. In the absence of internal signalling peptides that will guide them to the ER, proteins can follow alternative secretory pathways¹⁷ that can bypass the golgi and be transmitted directly to the cell membrane or be mediated via caspase-1¹⁷. Non-conventional secretory pathways can be determining for the protein's biochemical properties, e.g. FGF-2 follows ER/Golgi-independent secretion and when secreted via ER/golgi loses its ability to bind to HSPGs¹⁸.

1.1.3 Cell number regulation

Tissue cell number is a balance between cell proliferation and cell death. In tissue, stem- and precursor cells reside without proliferating until a demand for new cells arises. Hence, these cells undergo transition from G1 to S phase by surpassing the restriction points in G1. This transition is regulated by a 'gas and break' system of signalling. The 'gas' will lead the cells to proliferation, while the 'break' will trigger (1) cell cycle arrest where cells will remain in G1 phase (also called G0) until further stimulation, (2) senescence where the cells will lose their ability to enter the cell cycle but will remain metabolically active or (3) cell death.

The gas in the transition of cells from G1 to S phase is the presence of mitogens which stimulate expression/activation of proteins needed for entering S phase by surpassing the G1/S restriction point. Mitogens such as EGF, insulin-like growth factor (IGF) and platelet-derived growth

factor (PDGF) are responsible for the primary signal. Mitogens bind to receptor tyrosine kinases (RTKs) causing their dimerization.^{19,20} Activation of RTKs leads to phosphorylation and activation of Phosphoinositide 3-kinase (PIK3) that transiently activates Akt, thus causing inactivation of GSK3 β by phosphorylation²⁰. GSK3 β when active phosphorylates β -catenin that will lead to its degradation before translocation to the nucleus, activation of E2F and initiation of transcription²¹ (Figure 2). Additionally, GSK3 β can regulate degradation of cyclin D1^{22,23} that is vital for the transition from G1 to S phase²⁴. Apart from Akt²⁵, GSK3 β activity and β -catenin accumulation can also be regulated by the Wnt signalling pathway²⁶. Additionally after surpassing the restriction point controlled by Rb, transition from G1 to S phase requires the anchorage of cells which constitutes a second restriction point that is controlled independently of Rb²⁷.

Cell number is not only regulated by mitogens but also by cell cycle arrest, senescence and apoptosis signals that make up the 'break' in cell cycle progression. In the absence of mitogenic stimuli (e.g. in vitro during serum deprivation), cells will remain at the phase before the G1 restriction point (cell cycle arrest/G0) but will retain the ability to start proliferation and this phase is characterized by low levels of cyclins²⁸. However, cells can also go into hypermitogenic cell cycle arrest that, unlike G0, is characterized by high levels of cyclins and cyclin dependent kinases (CDKs) and activated mitogenic pathways but also the activation of CDK inhibitors such as p21 and p16²⁸. Due to the activation of CDK inhibitors, evasion of hypermitogenic cell cycle arrest can occur not by the presence of mitogens but instead by activation of downstream targets such as E2F²⁸. When ability for cell cycle progress is lost due to telomere shortage or by the presence of stress factor e.g. reactive oxygen species (ROS) or non-physiological conditions cells can undergo irreversible cell cycle arrest named senescence²⁸. The response and molecular mechanisms controlling senescence vary depending on the stimuli²⁹, but in most cases, cell senescence involves p53 activation which in turn leads to the activation of CDK inhibitors p16, p21 and p15^{29,30} that inhibit the phosphorylation of pRb thus leaving it active and able to suppress E2F³¹. Interestingly, unlike cells in cycle arrest, senescent cells remain metabolically active²⁹. Cell cycle progress is dependent on inactivation or down-regulation of apoptotic signals from anti-oncogenic proteins. For instance p53, p38, c-myc and JNK need to be down-regulated for cell cycle progress^{32,33}, however c-myc has been shown to promote apoptosis or proliferation depending on the concentration of mitogens and its activity has been shown to be regulated also via the Wnt signalling pathway³⁴. Apart from surpassing the restriction points, completion of cell cycle includes an increase in cell mass (cell growth). Mechanistic target of rapamycin (mTOR) is one of the main regulators of cell growth stimulation and its activity has also been associated with phosphorylation by Akt²⁰.




Figure 2. Cell cycle progression. Mitogens bind to RTKs leading to their activation. Downstream signalling activating PIK3 and Akt leads to de-activation of GSK3β and β-catenin translocation to the nucleus and transcription initiation. In the presence of high concentrations of mitogens, cells go into cell cycle arrest by expression of cdk inhibitors. Absence of mitogens leads to cell cycle arrest (G0). In the presence of stress or other stimuli cells lose their ability to enter the cell cycle and go into senescence controlled by p21, p16, p15 and p53. R= restriction point

1.1.4 Cell recruitment and migration

When there is need for tissue regeneration and /or tissue repair, recruitment of monocytes and stem cells from the blood stream to the tissue site is necessary. New cells need to migrate and infiltrate the tissue site where they proliferate and mature to differentiated cells in order to form new functional tissue. For recruitment, circulating cells in the blood need to be exposed and react to chemokines, adhere to and cross the endothelial barrier and finally move to the site where tissue needs to be repopulated.

Cells can be recruited in the presence or absence of an inflammatory signal. In the absence of inflammatory signals migration can be mediated via CXCL14³⁵. Inflammation -induced migration of monocytes is regulated via Chemokine (C-C motif) ligands (CCL) CCL2, CCL3 and Chemokine (C-X-C motif) ligand CX3CL1³⁵. Additional chemokines (e.g. CCL9) are responsible for their further recruitment to the lymph nodes. When chemokines bind to their receptors in the cells, they activate Ras-proximate-1 (RAP1) via cyclic adenosine monophosphate (cAMP) and exchange protein directly activated by cAMP (EPAC) which in turn binds to RALP. The RALP and RAP1 complex is associated to integrins and together with talin they induce morphological changes in the tails of α and β subunits of integrins giving them the ability to bind to intracellular adhesion molecular 1 (ICAM1) and vascular cell-adhesion molecule 1 (VCAM1) that are present on the surface of endothelial cells, thus inducing stronger adhesion. Integrins do not have activity potential on their own so they associate with other proteins and form complexes³⁶. The C-terminal tails of integrins associate with focal adhesion

kinase (FAK) and paxillin which cause the complex in a phosphorylation-dependent manner to bind to vinculin and actin via talin. This complex leads to actin reorganization with migration and adhesion effects^{36,37}. Moreover, activation of integrins leads to Rho GTPases activation that has an effect in actin re-organization. Integrin also causes up-regulation of lipid kinase activity and subsequent increase of Phosphatidylinositol 4,5-bisphosphate (PIP₂) that lead to downstream signalling of Akt, Erk and c-Jun N-terminal kinase (JNK) and thus cause effects on proliferation, survival and differentiation. Additionally, increase in PIP₃ leads to cytoskeletal re-organization and changes in cell polarity.

Stem cells are also recruited to the tissue when there is a need for tissue repair or regeneration. Stem cell recruitment and migration is also mediated via chemokine, chemokine receptor and integrin activation as described above. Chemokine Stromal cell-derived factor 1 (SDF-1) and its receptor CXCR4 are the main mediators of stem cell migration, and SDF-1 is expressed in bone marrow leading to stem cells remaining there. When there is a need for stem cell recruitment to tissue, this tissue releases SDF-1 leading to recruitment of stem cells to the site where they are needed by the same downstream pathway as described for monocytes³⁸.

1.1.5 Cell differentiation

Once precursor/progenitor cells have reached the site where tissue repair or tissue regeneration is required, exposure to the microenvironment leads to their cell differentiation. Factors in the microenvironment, growth factors and their respective receptors can have a determining effect on the lineage commitment of cells and their subsequent differentiation. Mesenchymal stem cell commitment to a specific lineage e.g. bone or adipose is regulated by various signalling proteins such as wnts³⁹, bone morphogenetic proteins (BMPs)⁴⁰, but also by constituents that comprise the extracellular matrix (ECM)⁴¹. Wnt signalling occurs via pathways known as the canonical pathway and an alternative pathway named non-canonical pathway (Figure 3). Wnts in the canonical pathway (Wnt10b) act by binding to low-density lipoprotein receptor-related protein (LRP) co-receptors and frizzled receptor leading to structural changes in frizzled causing the binding and activation of Dishevelled and inhibition of GSK3 β activity⁴². Canonical wnt signalling affects mesenchymal stem cells in various ways, resulting in inhibition or promotion of osteogenesis and adipogenesis depending on the level of maturation/differentiation of the mesenchymal stem cells³⁹.

Mesenchymal stem cells in early stages of their lineage differentiation remain proliferative and undifferentiated when canonical wnt signalling is active. In MSC committed to osteoblast lineage cells (pre-osteoblasts) but not mature cells, wnt10, wnt3a and wnt1 can promote differentiation by activating the transcription of Runx2¹¹. In committed adipocyte lineage cells (pre-adipocytes), canonical wnt signalling mainly by ligand wnt10b can inhibit differentiation by blocking CCAAT/enhancer binding protein α (C/EBP α) and PPAR γ and the downstream protein adipocyte protein 2 (aP2)⁴³.

Non-canonical wnt pathways act independently of the frizzled receptor and can also mediate β -catenin accumulation or act independently of β -catenin (Figure 3). Wnt5a binding to other receptors than frizzled, also inhibits adipogenesis, by activating SETB1 that forms complex with chromodomain-helicase-DNA-binding protein 7 (CHD7) and nemo-like kinase NLK preventing peroxisome proliferator-activated receptor α (PPAR γ) from transcribing adipocyte differentiation-related genes. wnt5b on the other hand, by binding to receptors other than frizzled, prevents β -catenin from translocating to the nucleus thus promoting adipocyte differentiation⁴³.




Figure 3. Wnt signalling pathways. A) Canonical Wnt signalling pathway. Wnt 10b binds to frizzled and LRP 5/6 leading to complex formation with dishevelled, inhibition of GSK3β and translocation of β-catenin to the nucleus. B) Non-canonical Wnt pathway, Wnt 5b binds to receptors that lead to inhibition of translocation of β-catenin to the nucleus and inhibition of TCF transcription. C) Non-canonical β-catenin independent pathway. Wnt 5a binds to receptors leading to complex formation of CHD7, SETDB1 and NLK and transcription of PPAR γ

1.2 ADIPOSE TISSUE HOMEOSTASIS

1.2.1 Adipose tissue as an endocrine and metabolic organ

Adipose tissue is a metabolic organ responsible for energy uptake, storage of triglycerides (TG) and release of free fatty acids (FFA) ⁴⁴, for hormone production and secretion and for secretion of adipokines that contribute to cross-talk with other organs and regulation of intermediary metabolism ⁴⁵. Hormones secreted from adipose tissue e.g. leptin have been shown to affect metabolism, whole body homeostasis and energy regulation ^{46–48}.

1.2.1.1 Cell types in adipose tissue

Adipose tissue comprises mainly of adipocytes, the differentiated tissue specific cells that store fat. Adipocytes are characterized by the presence of one or several lipid droplet(s) that occupy the majority of the cell cytoplasm and which are controlled in size by the presence of perilipin ⁴⁹. Adipose tissue additionally contains adipocyte precursors (mesenchymal stem cells and pre-adipocytes) that reside in the tissue until tissue regeneration is needed, when they mature to fully differentiated adipocytes and resident macrophages that are involved in tissue homeostasis.

1.2.1.2 Main components of the ECM in adipose tissue

The ECM composition of adipose tissue plays an important role in adipocyte maturation and is necessary for the change in the cell-cell contact and change of the cell morphology during adipocyte differentiation⁴⁴. During adipocyte differentiation, changes in the composition of ECM have been observed such as increases in collagen IV, laminin and nidogens⁵⁰⁻⁵³

1.2.1.3 Cell number

Formation of adipose tissue occurs from mesoderm⁵⁴ during early stages of development. Interestingly, even though the number of adipocytes remains stable during adulthood there is a 10% annual turnover^{55,56} of adipocytes, resulting to a need for adipocyte development from pre-adipocytes and mesenchymal stem cells. There are various different factors contributing to adipose tissue homeostasis and turnover, among which resident macrophages⁵⁷ and ECM components⁴⁴ appears to be of particular importance.

1.2.2 Mesenchymal stem cells

Adipocytes, the main cell type in adipose tissue are derived from mesenchymal stem cells. MSCs were first described by Friedstein et al. in 1966⁵⁸ and are non-hematopoietic stem cells that have the ability to differentiate into various types of tissue such as muscle, cartilage, bone and adipose tissue (Figure 4)^{59,60}.




Figure 4. Mesenchymal stem cell lineage. Mesenchymal stem cell differentiation can lead to several different types of tissue. Trans-differentiation (TD) potential has been observed between adipocytes and osteoblasts and between chondrocytes and osteoblasts indicating a close relationship between those pre-cursors

Among the cell types that MSCs can differentiate to, pre-osteoblasts and pre-adipocytes are very close phenotypically and depend on the same signalling pathways that could inhibit differentiation towards one lineage and activate the other ^{39,61,62}. Wnt signalling, BMPs and PPAR γ can guide the signalling towards adipogenesis or osteogenesis from MSCs, where e.g. wnt10b and wnt 5a lead the cells towards osteogenesis while wnt 5b leads the cells towards adipogenesis ² and it has also been shown that differentiated cells can transdifferentiate from one cell type to the other, e.g. osteoblast can trans-differentiate to adipocytes ^{63,64} but also chondrocytes can trans-differentiate to osteoblasts ^{65,66}.

1.2.3 From pre-adipocytes to adipocytes

Adipocyte differentiation has been studied extensively in vitro by the use of the murine fibroblast-like cell line 3T3-L1. In 1974, Meuth and Green described the ability of 3T3-L1 cells who have a fibroblast-like morphology to acquire lipid droplets and mature to adipocytes after stimulation with a hormone cocktail ⁶⁷ and since then 3T3-L1 have been used as a model for both cell cycle and differentiation studies of pre-adipocytes to adipocytes. Pre-adipocyte to mature adipocyte progress includes various stages such as growth arrest, clonal expansion, early differentiation and terminal differentiation (Figure 5).




Figure 5. Adipocyte differentiation stages. Pre-adipocytes go into clonal expansion followed by growth arrest and characterized by increased levels of C/EBP α and PPAR γ . Early differentiation is characterized by lipid acquisition, morphological changes, Akt signalling and Pref-1 down-regulation. Terminal differentiation is characterized by a merging of lipid droplets to one lipid droplet and expression of late markers e.g. GLUT-4.

1.2.3.1 Pre-adipocyte lineage

Characterization of cells as pre-adipocytes refers to cells that preserve the ability to proliferate, have a fibroblast-like morphology but are committed to the adipocyte cell lineage^{2,68}. The main difference of pre-adipocytes to other fibroblast like cells that derive from mesenchymal stem cells is the presence of pre-adipocyte factor-1 (pref-1) protein. pref-1 is expressed in pre-adipocytes and its down-regulation is required for the initiation of adipocyte differentiation⁶⁹. Pre-adipocytes reside in the adipose tissue and can be stimulated to proliferate and/or differentiate when it is needed and are retaining this capacity throughout life⁷⁰. Tissue expansion starts with proliferation of pre-adipocytes that is followed by their differentiation to mature adipocytes. Pre-adipocytes are the principal cells in the adipose tissue that have a proliferative capacity and ability to differentiate into mature adipocytes^{71,72}.

1.2.3.2 Pre-adipocyte cell number regulation

Regulation of pre-adipocyte number is dependent on the control of the cell cycle. As with other cell types, G1 to S phase transition is controlled by growth factors that stimulate the downstream expression of CDKs. Insulin-like growth factor have been shown to participate in the stimulation of proliferation of pre-adipocytes⁷³ and also to induce resistance to stress factors and apoptosis in these cells. Insulin-like growth factor- binding proteins have also been reported to affect pre-adipocyte proliferation⁷⁴. Finally hormones such as Thyroxines (T3, T4), glukocortikoids, adrenalin, noradrenalin have an impact on the metabolism of adipose tissue (lipogenesis and lipolysis).

1.2.3.3 Growth arrest

Once cells have reached confluence, cell-cell contact leads to cell cycle arrest. This process, however, has been shown not to be required for the differentiation of pre-adipocytes in vitro⁴⁴. C/EBP α has been shown to regulate the cell growth arrest of pre-adipocytes by p21 mRNA and protein expression^{75,76}. Transcription factor PPAR γ can also induce cell growth arrest by reducing the DNA binding capacity of E2F/DP⁷⁷.

1.2.3.4 Clonal expansion

When differentiation is initiated, cell growth- arrested pre-adipocytes need to re-enter the cell cycle, at which stage they undergo proliferation known as clonal expansion. Interestingly it has been suggested that the molecular pathways for cell cycle proliferation before and after cell growth arrest (clonal expansion) differ⁴⁴. Post- confluent pre-adipocyte clonal expansion is mediated via changes in the phosphorylation status of Rb and the transient increase of p107⁷⁸. However, cell cycle arrest that is required before early differentiation and is controlled by the expression of C/EBP⁷² and is not mediated by Rb and Cdk inhibitors but can take place even when Rb is hypophosphorylated⁷⁹.

1.2.3.5 Early differentiation

Early differentiation stage of adipogenesis is characterized by the up-regulation of several transcription factors. Main regulators of adipocyte differentiation C/EBP and PPAR γ are dramatically increasing during early differentiation stages. Moreover, the anti-adipogenic factor that is specific for pre-adipocytes, pref-1, is down-regulated at this stage. During early differentiation, cell morphology and ECM composition undergo important changes. More specifically, actin and tubulin are reduced⁸⁰, while there is a switch in the collagen types with an increase of collagen type IV and a decrease of type I and III procollagen and an increase in nidogen expression^{51,81}. Akt signalling is essential for adipogenesis^{82,83} and Akt knock-down

impairs expression of Krüppel-like factor that is required for completion of adipocyte differentiation^{84,85}.

1.2.3.6 Terminal differentiation

The final stages of adipocyte differentiation are characterized by rapid accumulation of lipids, mainly TG, fusion of small lipid droplets in to one large lipid vacuole that covers most of the cytoplasm of the cell. Terminal differentiation is also accompanied with increased insulin sensitivity and expression of perilipin, Glucose transporter type 4 (GLUT-4) and enzymes involved in triacylglycerol metabolism⁴⁴. ECM re-organization has been shown to be essential for terminal differentiation and it has been shown that inhibition of collagen production impairs terminal adipocyte differentiation⁸⁶.

1.2.4 Macrophages of adipose tissue

1.2.4.1 Types of macrophages in adipose tissue

In adipose tissue the majority of immune cells are macrophages, also known as adipose tissue macrophages (ATMs). In lean adipose tissue the majority of ATMs are anti-inflammatory⁸⁷. ATMs are generally classified into two groups, the pro-inflammatory cytokine secreting group M1 that is characterized by secretion of e.g. tumor necrosis factor α (TNF- α) that participate in the tissues immune defence and the anti-inflammatory cytokine secreting group M2 that secrete e.g. IL-4 that are regulating tissue homeostasis and can take part in tissue repair and remodelling^{88,89}. ATMs can be recruited to the site by adipokines secreted from adipocytes⁹⁰ but it has also been shown that resident macrophages are able to proliferate by stimulation with IL-4 and do not need to be recruited from the blood stream⁹¹. In obesity, the ratio between anti-inflammatory (M2) and proinflammatory (M1) macrophages changes towards the majority of macrophages being polarized into M1 pro-inflammatory and this switch is triggered by secretion of adipokines from adipocytes as well as from an autocrine system from macrophages themselves^{88,92-94}.

1.2.4.2 Anti-inflammatory macrophages (M2)

Anti-inflammatory macrophages (M2) that reside in healthy adipose tissue, are activated by PPAR γ and are positively regulating insulin-sensitivity⁹⁵. Contrary to the popular belief that macrophages are hindering adipocyte differentiation and development it has also been shown that impairment of pro-inflammatory activation in mice leads to reduced capacity of tissue expansion⁵⁷ and it is known that M2 macrophages increase anti-inflammatory cytokines like IL-10 and participate in tissue repair⁸⁹ M2 macrophages can trigger tissue repair by the production of MMPs leading to ECM remodelling, growth factors and cytokines that trigger proliferation and differentiation of precursor cells in tissue⁹⁶ and are responsible for metabolic effects such as protection from insulin resistance via production of IL-10^{92,95}.

1.2.4.3 Pro-inflammatory macrophages (M1)

On the other hand, M1 pro-inflammatory macrophages that are recruited and triggered by the expression of TNF- α , monocyte chemoattractant protein-1 (MCP-1) and adipokines are correlated to BMI and have been shown to contribute to the metabolic syndrome symptoms, tissue remodelling, and chronic inflammation^{88,89,93,94,97,98}.

1.2.5 Adipose tissue changes in obesity

Adipose tissue is now considered an important secretory organ for adipokines that influences the function of several organs ⁹⁹, thus changes in tissue homeostasis or adipose tissue malfunctions leads to systemic problems. Obesity has been shown to be associated with an increase in the amount of ATMs ¹⁰⁰ presumably due to increased death of adipocytes ¹⁰¹ where macrophages gather to remove the dead cells and hypoxia in the tissue ¹⁰² leading to chronic low grade inflammation ¹⁰³. Inflammation in adipose tissue has been shown as the cause of several conditions such as cardiovascular disease, atherosclerosis, type-2 diabetes and dyslipidemia that all together are referred to as the metabolic syndrome ¹⁰⁴.

Adipose tissue expansion results in secretion of adipokines that will trigger the recruitment of macrophages and the 'phenotypic switch' of macrophages from anti-inflammatory M2 to pro-inflammatory M1 macrophages. Increase in pro-inflammatory M1 macrophages is mainly regulated by the MCP-1/CCR2 signalling that causes the recruitment of macrophages to adipose tissue ^{105,106}. MCP-1 and free fatty acid release (caused by lipolysis) are secreted by adipocytes that have become hypertrophic thus triggering the signalling for M1 macrophages ¹⁰⁷ (Figure 6).

Obesity in adults is mainly characterized by the enlargement of the already existing adipocytes (hypertrophic obesity) and in some cases proliferation and increase in the adipocyte number (hyperplastic obesity) ¹⁰⁸. Hyperplasia is associated with better insulin response and glucose metabolism compared to hypertrophy in obesity ¹⁰⁹. Proliferation of pre-adipocytes occurs only after the full expansion of adipocytes ¹¹⁰ and cell growth of adipocytes leads to up-regulation and release of growth factors that can both promote the proliferation of pre-adipocytes e.g. IGF-I ¹¹¹.




Figure 6. Adipose tissue remodelling in obese state. Increase of adipose tissue is characterized by increased TNF α , free fatty acids and hypoxia. These changes lead to macrophage recruitment and macrophage proliferation and morphological switch from anti-inflammatory m2 macrophages to pro-inflammatory M1 macrophages.

1.3 TARTRATE- RESISTANT ACID PHOSPHATASE

1.3.1 TRAP history and properties

Tartrate- resistant acid phosphatase (TRAP; E.C. 3.1.3.2), also known as purple acid phosphatase (PAP), Acp 5 or uteroferrin (Uf) ^{112,113} is a diiron centre- containing metalloenzyme ^{114,115}. TRAP was initially described as a member of acid phosphatases in 1954 by Sundararajan and Sarma ¹¹⁶ and acid phosphatase properties were investigated extensively in 1970s ^{117,118}. In 1971, Yam et al. described an isoenzyme of acid phosphatase that was tartrate- resistant and highly expressed in cells from the rare B-lymphocyte-derived hairy cell leukaemia ¹¹⁹. During the 1970s and 1980s a variety of studies described common properties for purple acid phosphatase and tartrate- resistant acid phosphatase and in 1990s a study from Vincent and Averill proposed that purple acid phosphatase and tartrate resistant acid phosphatase were identical ¹¹³. TRAP has traditionally been used as a marker for osteoclasts and activated macrophages ^{120–122}, however in recent years more functions for TRAP have emerged.

1.3.1.1 TRAP gene and translation

The gene encoding TRAP (Acp5) is located on chromosome 19p13.2-13.3 in humans ^{123,124} and chromosome 9 in mouse ¹²⁵ and contains five exons of which 4 are translated into protein sequences ^{126,127} while the untranslated exon 1 exists in 3 different variants in mouse (4 in humans) with giving rise to 3 mRNA transcripts encoding identical protein sequences but with differential cell- and tissue expression patterns ^{128–130}.

1.3.1.2 Structure-proteolytic cleavage

TRAP exists in two different isoforms named TRAP 5a and TRAP 5b ^{131,132}. The product from translation of the Acp5 gene is the 35 kDa TRAP isoform denoted 5a that can undergo proteolytic cleavage giving rise to the dimeric isoform TRAP 5b of ~24 and 16 kDa “subunits”, held together by a disulphide bond ¹³³ (Figure 7). Proteolytic cleavage of TRAP 5a occurs in a repressive loop region, Ser145-Val161 ^{115,133} by various proteases, such as cathepsins, trypsin and papain and is responsible for the increased phosphatase activity of TRAP 5b ^{133–136}. The kinetic properties of TRAP are additionally influenced by the two N-linked oligosaccharide chains present in the protein ¹³⁷.

Despite the fact that for many years isoform TRAP 5a was considered to serve only as the inactive precursor to the enzymatically active isoform TRAP5b, it has recently been apparent from several studies that the two isoforms can indeed have different functions and localizations and that TRAP 5a has a separate function promoting pre-adipocyte proliferation and differentiation and when over-expressed in mice leads to hyperplastic insulin sensitive obesity ¹³⁸.




Figure 7. Proteolytic cleavage of TRAP 5a generates TRAP 5b. TRAP 5a is cleaved by proteases in a repressive loop that generates the “dimer” TRAP 5a held together by disulphide bonds and leading to changes in the di-iron centre increasing the phosphatase activity of the protein.

1.3.2 TRAP expression and localization

1.3.2.1 TRAP in macrophages and dendritic cells

TRAP is expressed by a subset of macrophages^{139 140} and dendritic cells¹⁴¹ and correlates to their activation or maturational state. When TRAP is genetically knocked down in mice, an increased basal expression of inflammatory cytokines such as IL-1b, IL-12 and TNF α was noted¹⁴² while dendritic cells in mice lacking TRAP have increased expression of IL-10 but reduced response to T- dependent antigens¹⁴³. The expression of TRAP is also regulated by cytokines and it has been observed that TRAP is reduced in response to IFN γ and LPS¹⁴⁴ but another study indicated increased TRAP expression by LPS¹⁴⁵ and additionally TRAP expression is increased by TNF α in monocytes¹⁴⁶

1.3.2.2 TRAP in bone

In bone TRAP has been shown to be expressed and secreted by osteoclasts. More specifically, TRAP 5a is secreted to the extracellular matrix where it is subjected to proteolytic cleavage giving rise to TRAP 5b while intracellular TRAP is found as both isoforms TRAP 5a and TRAP 5b^{133,147,148}.

Apart from osteoclasts, TRAP in bone is expressed by osteoblasts and osteocytes and its expression is independent of the osteoclast activity^{149–153} but is increased in rats with osteoporosis¹⁵⁴. Additionally, when TRAP is over-expressed in mice there is increased bone mineral content and density that stems from increased osteoblast differentiation since resorption activity remains stable¹⁵⁵ indicating a function of osteoblast TRAP different than that of osteoclast TRAP. TRAP in osteoblasts is located in LAMP-1 positive vesicles thus exhibiting lysosomal localization¹⁵⁶.

Finally TRAP has been shown to be expressed in cartilage by hypertrophic chondrocytes¹⁵⁶. This may explain the cartilage phenotype observed in TRAP-deficient mice where the growth plate displayed increased height and disorganized columns of chondrocytes¹⁵⁷. TRAP expression from chondrocytes and osteoblasts is of interest since they share a common mesenchymal stem cell lineage and particularly since TRAP expression in cells of the same lineage such as pre-adipocytes is very low¹⁵⁸, indicating a regulation of TRAP expression potential during commitment to different lineages.

1.3.2.3 TRAP expression from other cell types

Apart from the immune system and bone where TRAP expression has been extensively studied, TRAP expression has been observed in other cell types such as neurons¹⁵⁹, hepatocytes and keratinocytes¹⁶⁰.

1.3.2.4 TRAP binding partners

TRAP has been shown to interact with heparin^{148,161,162} and heparin has been shown to inhibit activity of TRAP 5a but not TRAP 5b¹⁶³. TRAP has also been suggested to interact with the HPSG glypican-4 in bone¹⁶⁴ and with the TGF β receptor interacting protein 1 (TRIP-1)^{152,165,166}. In serum, TRAP has been shown to be associated to α 2-macroglobulin in a calcium-dependent manner¹⁶⁷.

1.3.3 TRAP in disease

1.3.3.1 TRAP 5a vs. TRAP 5b as disease markers

TRAP 5a and TRAP 5b have been suggested to have different localizations and serve as markers to different pathological conditions, where TRAP 5b has been shown to be a marker for osteoclast number and bone resorption in serum^{133,120–122} and end-stage renal disease¹⁶⁸ and kidney failure. On the other hand TRAP 5a has been suggested as a marker for mostly inflammatory conditions¹⁶⁹, such as atherosclerosis¹⁷⁰, rheumatoid arthritis,^{171,172} and sarcoidosis¹⁷³.

1.3.3.2 TRAP and autoimmunity

Besides expression TRAP in certain macrophages, also certain dendritic cells known to be involved in antigen presentation, have been shown to express TRAP with a proposed role in dendritic cell maturation¹⁴². Knock-down of TRAP in mice and in humans with inactivating mutations in the TRAP gene show a similar deregulated immune phenotype with increased basal expression of Th1 cytokines such as Interferon- γ (IFN γ), TNF α and IL-1^{142,174}. This aberrant immune response was coupled to increased phosphorylation of osteopontin, a protein with known function to control TH1/Th2 bias through dephosphorylation by TRAP^{175–177}.

1.3.3.3 TRAP in cancer

TRAP has also been shown to be elevated in melanoma, gastric, colon, ovarian and breast cancer^{178–183} and when over-expressed in a human breast cancer cell line, TRAP 5a and TRAP 5b can be found intracellularly while only TRAP 5a is secreted¹⁸⁴. Moreover, TRAP has been shown to increase invasion and metastasis of cancer cells, to increase their migration and change their morphology and adhesion pattern¹⁸⁵. In contrast, siRNA transfection revealed a morphological effect consistent with impaired detachment to plastic-bound substrates¹⁸⁶.

1.3.4 Regulation of cell number and differentiation by TRAP

TRAP has been shown to have effects on cell proliferation and differentiation^{152,155}. In bone tissue TRAP has been shown to give rise to increased cortical bone mineral density and content and increased expression of bone differentiation markers when over-expressed¹⁵⁵ in mice and to have an effect on differentiation of osteoblasts^{165,187} and to interact with the TGF- β -receptor-interacting protein (TRIP-1) that exerts a TGF- β like differentiation^{152,166}. Osteoblast-like cells have also exhibited the ability to both express and endocytose TRAP^{151,153} and TRAP levels are altered during osteoblast differentiation¹⁸⁷. However, the exact function of TRAP expression and endocytosis by osteoblast-like cells in bone is not known. TRAP has additionally been shown to have an effect on cell number regulation of hematopoietic cells and to be a regulator of fetal erythropoiesis by acting synergistically with erythropoietin and other growth factors^{188,189}.

1.3.5 Regulation of migration by TRAP

Apart from cell number regulation TRAP has been shown to affect the motility, morphology and migration of cancer cells¹⁸⁵. In osteoclasts, TRAP exhibits phosphatase activity with osteopontin as substrate^{175,176} and dephosphorylation of osteopontin leads to reduced attachment of osteoclasts and changes in migration^{190,191}.

1.3.6 TRAP 5a in adipose tissue

Interestingly, isoform TRAP 5a has been shown to be involved in the regulation of the adipose tissue. In 2008, Lång et al. showed that when TRAP 5a is overexpressed in mice, it leads to hyperplastic insulin- sensitive obesity and that human and mouse pre-adipocytes show increased proliferation and differentiation by the addition of TRAP 5a but not 5b¹³⁸. Additionally, no signs of inflammation were observed in the TRAP 5a over-expressing mice¹³⁸.

In adipose tissue, TRAP 5a has been shown to be elevated in obese individuals¹⁹² and an increase of TRAP 5a has been described in the serum of overweight young males¹⁹³. However, unlike other inflammatory markers, its expression was not reduced after exercise and weight loss¹⁹³. TRAP expression in different adipose tissue depots also changes with obesity. In lean individuals TRAP levels are higher in visceral, while in overweight and obese individuals, higher levels of TRAP are observed in subcutaneous fat depot⁹⁸. TRAP 5a expression and secretion in the adipose tissue is almost exclusively from macrophages^{138,192} and even though TRAP 5a and TRAP 5b are both found intracellularly in ATMs, TRAP 5a is the only isoform that gets secreted¹⁹². Additionally, TRAP levels co-related to both number and size of adipocytes as well as with TNF α and IL-6¹⁹².

TRAP 5a secretion by macrophage suggests a paracrine action where TRAP 5a is secreted by macrophages but can act on pre-adipocytes and affect their proliferation and differentiation. However, until now little is known on the mechanism of action of TRAP 5a on cells in order to regulate cell number, differentiation and migration as well as its binding partners on the extracellular environment and cells.

1.3.7 Working hypothesis

The working hypothesis of this thesis was that TRAP 5a is secreted by a subset of macrophages in the adipose tissue, where it is presented to pre-adipocytes for binding to the cell surface and initiation of signalling pathways that can promote proliferation, differentiation and further effects in pre-adipocytes (Figure 8). However the molecular pathways, binding partners and secretion patterns of TRAP 5a in adipose tissue remain to be explained. In this thesis, the aim was to shed light on different aspects of TRAP 5a action in adipose tissue such as expression levels related to BMI, interaction of TRAP 5a with binding partners in ECM and cell surface, signalling pathways and functional effects in pre-adipocytes.




Figure 8. Working hypothesis of TRAP 5a mechanism of action in adipose tissue. TRAP 5a secreted from macrophages acts on pre-adipocytes in order to stimulate proliferation and differentiation as well as possible unknown effects.

2 AIMS OF THE THESIS

Paper I. The aim was to develop a human TRAP 5a specific ELISA in order to evaluate levels of TRAP 5a from obese vs. lean individuals.

Paper II. The aim was to identify potential binding partners of TRAP 5a in adipose tissue.

Paper III. The aim was to investigate the interaction between TRAP 5a and pre-adipocytes, the binding mechanism on the cell surface, endocytosis routes and interaction with other cell types of mesenchymal lineage.

Paper IV. The aim was to investigate the effect of TRAP 5a in the regulation of pre-adipocyte cell number, differentiation and migration and identify potential signalling pathways through which TRAP 5a action occurs.

3 REMARKS ON METHODOLOGY

3.1 TRAP 5A ELISA DEVELOPMENT

For ELISA development, the first step was the generation of monoclonal antibodies specific for human TRAP 5a. To achieve this, the peptide loop region present on human TRAP 5a but not human TRAP 5b was used as the immunogen by which mice were immunized. Apart from TRAP 5a antibodies, antibodies against total TRAP (TRAP 5a+TRAP 5b) were developed. To achieve this, the full length protein was used. Next, isolated antibodies were epitope mapped by ELISA. Epitope mapping of antibodies allows the grouping of antibodies for further use in a sandwich ELISA. The use of antibodies from different epitopes increases the possibility of both capture and detection antibodies to bind to the protein with high affinity and no interference of binding from one to the other.

One aspect of the quality assessment of the newly developed ELISA was the effect that serum components can possibly have on the detection of TRAP 5a. For this purpose recombinant TRAP 5a was diluted in serum in a dose- dependent manner and absorbance was measured.

The sandwich ELISA was evaluated by measuring absorbance in a TRAP 5a dose response, testing of TRAP 5b absorbance and measuring intra- and inter- assay variation and reference interval in a set of human serum samples from healthy males to verify the reproducibility of the results.

3.2 PULL DOWN ASSAYS

In paper II, the potential binding of TRAP 5a to nidogen-2 and TRIP-1 were studied in vitro using recombinant proteins in pull- down assays. In this system, His-tagged proteins were used as bait. The property of the His-tag to bind to Ni was used to bind the ‘bait’ proteins in magnetic agarose- Ni beads that could then be pulled-down by magnet. Ni- beads often exhibit considerable unspecific binding of proteins and a way to address this is saturation of beads with His-peptides to be used as control. In the pull- down assays described, control beads (beads with no His-tagged recombinant protein) were saturated with His- peptides to prevent unspecific binding.

One other important aspect to be taken into account when performing pull- down assays was the stoichiometry between the recombinant proteins that are studied. In the pull-down assays between His-tagged nidogen-2 domains and TRAP 5a, stoichiometry was always 1:1 for ‘bait’ and protein of interest.

3.3 ICC/ 2-STEP INDIRECT STAINING

To investigate if TRAP 5a can bind to the surface of pre-adipocytes, immunocytochemistry and indirect staining was used, where cells were treated with TRAP 5a for 1h and then fixed with formalin or acetone. After fixation cells were incubated with primary antibody against TRAP and fluorescent secondary antibody. The staining observed reflected only bound TRAP 5a since no TRAP 5a was detected in untreated cells. Since the potential binding partner for TRAP 5a was unidentified, to study binding of TRAP 5a cells were treated with TRAP 5a at 4°C to minimize endocytosis, thus ensuring that TRAP staining will only reflect TRAP 5a

bound to the surface. For the same purpose immunocytochemistry was performed without cell membrane permeabilization. Immunocytochemistry was followed by image analysis.

3.4 IMAGE ANALYSIS

In paper II, immunocytochemistry, proximity ligation assay and immunohistochemistry was used to evaluate nidogen-2 expression, interaction of TRAP 5a with nidogen-2 and TRIP-1 and co-localization of TRAP 5a with nidogen-2 in adipose tissue, respectively.

Image analysis was performed as follows: For proximity ligation assays the number of positive spots (each spots corresponds to an interaction site) were counted using Volocity software and divided with the area of each cell. For immunohistochemistry, co-localization was also quantified and analysed. In this case, co-localization was measured using Volocity software to calculate Pearson's correlation between the two colours (green for TRAP 5a and red for nidogen-2) as well as overlap coefficient M1 and M2 for the two channels, in five regions of interest. An important aspect of image analysis in order to generate Pearson's correlation and M1 and M2 values is setting threshold for signal. In all image analysis threshold was set by negative control of one of the two proteins, in order to minimize false positive co-localization. In adipose tissue, section that have not been stained for nidogen-2 served as negative control for threshold setting.

Overlap co-efficient refers to the amount of overlap of one channel towards the other, for instance M1 overlap coefficient refers to the amount of overlap of green channel to the red channel, while M2 refers to the overlap of red channel to the green channel. This is important in order to identify if two proteins are always co-localized (high M1 and M2 values) or if only one protein is always co-localized to the other (high M1 value but low M2 or vice versa) but the other one can be also observed in other areas.

Image analysis was also performed in immunocytochemistry and co-localization studies (paper III) following the same principles as described above. Since these studies were performed using pre-adipocytes, threshold was set for pre-adipocytes that had not been treated with TRAP 5a. Pre-adipocytes do not express TRAP 5a thus if not treated with TRAP 5a, no staining should be observed. This property can thus serve as a negative control. In this paper, quantification of ICC was also performed using velocity software and quantifying the number of clusters/area. For the quantification of clusters, parameters such as circularity, size and intensity of spots/clusters were standardized using TRAP 5a treated cells as a positive control. Data collected in the above method were statistically analysed in 4-10 cells using non-parametric Mann-Whitney U-test.

3.5 CELL CYCLE ANALYSIS

In paper IV, proliferation was studied by cell cycle analysis using the Bromodeoxyuridine (BrdU) incorporation method in combination with FACS analysis. This was performed in cells that were allowed to enter the cell cycle and left for various time points to progress in the cell cycle before BrdU was added. BrdU analysis can be performed in two manners (1) BrdU is added immediately after cell cycle entry and is incorporated into the DNA for several hours before cell fixation, staining and FACS analysis or (2) BrdU is added at various time points only for a short period of time before cell fixation. In the first case, BrdU incorporation is a measurement of the percentage of cells that have engaged in cell cycle entry immediately after serum stimulation and until the time of fixation, thus giving an overview of the total number of

cells undergoing cell cycle progress. On the other hand, in approach (2), BrdU incorporation occurs briefly and only to cells that undergo S phase at the moment of BrdU addition and not the total amount of cells that have undergone G1 to S transition. In this study the second approach was used where BrdU incorporation reflects the number of cells in S phase momentarily and throughout the whole time course. This approach was selected in order to evaluate at which time point TRAP 5a effect takes place without masking the effect from the accumulation of cells in the S phase.

The selection of BrdU for studying proliferation was based on two reasons. Firstly, BrdU incorporation in combination with propidium iodide (PI) can give a clear separation of the cell cycle stages G1, S and G2/M thus allowing for a detailed comprehension of the cell cycle and TRAP 5a effects. Second, viability assays broadly used for proliferation studies (e.g. XTT assay) are based on mitochondrial activity. Since TRAP 5a effects on cells are under investigation, it cannot be excluded that TRAP 5a alters the mitochondrial activity of the cells thus affecting the read-out of assays based on this type of activity.

3.6 PHOSPHORYLATION SIGNALLING ASSAYS

In paper IV, in order to investigate signaling assays triggered by TRAP 5a upon cell cycle entry, a signalling array detecting alterations in phosphorylation status was used. Since phosphorylation status of proteins involved in the cell cycle regulation occur very fast, the time points selected ranged between 5-30 min.

4 RESULTS AND DISCUSSION

4.1 PAPER I

In this study the aim was to develop a human TRAP isoform 5a- specific ELISA that would give the possibility of studying TRAP 5a levels in serum. Here, obese vs. lean individuals was compared in order to assess if TRAP 5a correlates to BMI in serum as previously reported ¹⁹³.

Firstly we developed monoclonal antibodies (mAbs) specific for TRAP 5a using the loop present in TRAP 5a but absent in TRAP 5b due to proteolytic cleavage ¹³¹ as well as mAbs detecting both isoforms using the full length protein. Epitope mapping allowed grouping the mAbs according to the recognition epitope. Three groups of mAbs were identified for total TRAP (TRAP 5a+5b) one of which was mAb with the peptide loop as recognition epitope (mAb 46). Results showed that TRAP 5a could successfully be recognized by the specific mAb 46 both in Western immunoblot and ELISA and that a mAb (clone 12.56) raised against full length TRAP could recognise both isoforms. With the combination of a mAb against TRAP 5a (mAb 46) as capture antibody and a mAb against total TRAP (mAb 12.56) as detection antibody, the development of an ELISA was possible. Even though, antibodies specific for TRAP 5a have been developed and used in the past ^{194,195}, this is the first time that mAb against TRAP 5a has been raised using the peptide loop present only in TRAP 5a, instead of subcloning which could potentially increase the specificity of an ELISA since it is assured that the recognition epitope is only present on isoform TRAP 5a.

After evaluation of mAbs, the ELISA was validated for isoform specificity using TRAP 5a and TRAP 5b isoforms and it was shown that with the combination of antibodies mAb 46 (TRAP 5a) and mAb 12.56 (TRAP 5a+5b) only TRAP 5a was detected and absorbance was dose-dependent with a lower detection limit of $0.112 \pm 0.039 \pm$ ng/ml. Since TRAP has previously been shown to associate with α 2-macroglobulin in serum ¹⁶⁷, the effect of serum in the dose-dependent response for TRAP 5a was evaluated by the use of commercially available serum samples. Hence it was shown that serum has no effect on TRAP concentrations up to 0.5ng/ml in the standard curve. At higher TRAP 5a concentrations, serum had an inhibitory effect on the absorbance which could be due to failure to completely dissociate complexes of TRAP 5a with serum components.

Inter- and intra-assay variation of TRAP were evaluated in a set of commercially available serum samples (3 samples for intra-assay and 17 samples for intra-assay) from healthy males. Intra-assay variation was calculated in average to 10.3% and inter-assay variation to 10.7% that are both in acceptable ranges for an ELISA assay. Apart from variations, the reference interval was calculated to 0.86-6.27 ng/ml while the mean concentration was 3.65 ± 1.23 ng/ml. The mean concentration for healthy males is in the same range as reported previous studies where TRAP 5a was measured and have shown that mean TRAP 5a concentration in serum of healthy males was 2.65 ± 0.29 ng/ml ¹⁹³ and 5.89 ± 1.65 ng/ml ¹⁹⁵ in different groups of healthy males.

After validation of the newly developed ELISA, TRAP 5a ELISA was used to measure the levels of TRAP 5a in serum from lean vs. obese females. The same groups were used to measure total TRAP with a previously described ELISA ¹⁹² in order to evaluate if the relationship between the groups is the same for TRAP 5a and total TRAP. The mean concentration for TRAP 5a of lean individuals was calculated to 3.78 ng/ml and to 1.97 ng/ml for obese females using the newly developed TRAP 5a ELISA indicating that TRAP 5a levels were lower in obese individuals compared to lean. These findings seem to be in contrast with what has been observed in a previous study where TRAP 5a was elevated with increased BMI ¹⁹³. However in this previous study the group investigated was overweight and male rather than obese (BMI=25-30 kg/m²) while in this study all individuals in the female obese group had a

BMI of over 30 kg/m². The same relation between lean and obese group was also indicated when total TRAP was measured, where the mean concentration for the lean group was 1.19 ng/ml and for the obese group 0.79 ng/ml. Despite the fact the two assays seem to decline in terms of absolute values, they seem to exhibit the same relation for TRAP levels between lean and obese group.

Finally, it was investigated whether there is a correlation between TRAP 5a and total TRAP levels with BMI for the group of lean and obese individuals. A significant correlation was only observed between TRAP 5a and BMI in the group of obese individuals, thus indicating that in lean individuals, multiple factors apart from the adipose tissue burden can contribute to the levels of TRAP 5a. In the study by Shih et al. in 2010¹⁹³, correlation was observed in a group of individuals with BMI of 18.5-30 kg/m² however no separation between lean and overweight group was performed.

In summary, in paper I the development and validation of a new ELISA for detection of TRAP 5a isoform is described, with the use of monoclonal antibodies raised against the peptide loop present in TRAP 5a but not TRAP 5b (for TRAP 5a specific mAb) or the full length protein (for total TRAP mAbs), its evaluation using serum from healthy males and its use to detect TRAP 5a levels in serum from obese and lean females. Results showed a decreased TRAP 5a expression in the serum of obese (BMI \geq 30 kg/m²) vs. lean females and correlation between TRAP 5a and BMI only in the obese group, perhaps indicating that only in obesity TRAP 5a levels are mainly determined by adipose tissue. It has also been shown that expansion and morphology of adipocytes in different depots (hyperplasia vs. hypertrophy) plays a role in the metabolic profile¹⁰⁹ and also that TRAP highest expression shifts from visceral in lean to subcutaneous in obese individuals⁹⁸, thus indicating that the depot and morphology of adipose tissue could also influence the serum levels.

4.2 PAPER II

The aim of this study was to identify potential binding partners for TRAP 5a in the adipose tissue and to investigate if the previously suggested binding partner for TRAP, i.e. TRIP-1^{152,165} is interacting with TRAP 5a in pre-adipocytes. For this purpose, cell membrane proteins from pre-adipocytes were isolated and used in a pull-down assay where His-tagged TRAP 5a was used as 'bait'.

After pull-down assay, 3 bands with molecular weight higher than TRAP 5a were observed by SDS-PAGE and silver staining and were analysed using mass spectrometry. All three bands contained a peptide sequence corresponding to protein nidogen-2. This finding was initially surprising since nidogen-2 is not known to be found on the membrane of cells, however, this could be due to nidogen-2 attaching to the cell surface via its RGD-region¹⁹⁶ and thus being isolated together with membrane proteins. Nidogen-2 has originally been shown to be an ECM protein and to have a high homology with nidogen-1¹⁹⁷ while its expression is up-regulated during adipocyte differentiation⁵², thus it was hypothesized that nidogen-2 could bind TRAP 5a after secretion of the latter from macrophages in the ECM in an interaction module similar to what has been observed for growth factors e.g. for HGF interaction with vitronectin and fibronectin¹⁹⁸.

Firstly, nidogen-2 mRNA and protein expression in pre-adipocytes and differentiating pre-adipocytes was investigated and it was shown that indeed mRNA expression of nidogen-2 is up-regulated during differentiation. Nidogen-2 was detected both in media and cell lysate of pre-adipocytes, however a large fraction was of lower molecular weight than full length nidogen-2, indicating a proteolytical processing event in one of its cleavage sites¹⁹⁹.

Localization of nidogen-2 in pre-adipocytes was investigated by immunocytochemistry and this protein was shown to be located in cytoplasmic vesicles and fiber-like structures, while in previous studies nidogen-2 was observed pericellularly ¹⁹⁹. Since identification of nidogen-2 by mass spectrometry had very low coverage, additional methods were used to verify if indeed it interacts with TRAP 5a. Pull down-assays were used with either His-tagged TRAP 5a or His-tagged nidogen-2 as 'baits' and untagged nidogen-2 or untagged TRAP 5a respectively. Nidogen-2 contains three distinct domains G1, G2 and rod-G3 with different binding motifs ¹⁹⁷ and pull-down assays with the different domains were also used to evaluate which domain of nidogen-2 mediates the binding to TRAP 5a. Results showed that TRAP 5a could pull down nidogen-2 and that nidogen-2 rod G3 domain was responsible for the interaction between TRAP 5a and nidogen-2, thus further verifying that the potential interaction between the two proteins. Since nidogen-2 was shown to be localized to vesicle- and fibre-like structures in pre-adipocytes, it was investigated whether there existed a potential interaction of nidogen-2 with TRAP 5a. For this purpose, proximity ligation assay was performed in pre-adipocytes that had been treated with TRAP 5a, showing that TRAP 5a and nidogen-2 are indeed interacting in pre-adipocytes. The interaction between nidogen-2 and TRAP 5a could occur in endocytic vesicles by endocytosis of TRAP 5a from the ECM, in complex with nidogen-2 fragments. Finally, the localization of TRAP 5a in relation to nidogen-2 in adipose tissue was investigated and it was shown that the two proteins are co-localized in vesicles in adipose tissue cells, indicating a potential interaction.

Apart from the interaction of nidogen-2 and TRAP 5a, the previously suggested binding partner for TRAP 5a, i.e. TRIP-1 was investigated for interaction with TRAP 5a in pre-adipocytes. This showed that TRIP-1 and TRAP 5a interact in vitro in a pull-down assay system and intracellularly in pre-adipocytes. Interaction between the two proteins intracellularly in vesicles was surprising since TRIP-1 has been suggested to be localized in the cytoplasmic matrix and nucleus ²⁰⁰ while TRAP 5a is not expressed by pre-adipocytes but only endocytosed and thus most likely located in endocytic vesicles.

In summary, this study aimed to identify new binding partners for TRAP 5a in pre-adipocytes and to investigate the previously suggested interaction of TRAP with TRIP-1 with the use of pull down and proximity ligation assays. Results revealed nidogen-2 as a potential binding partner of TRAP 5a intracellularly in pre-adipocytes and in adipose tissue and indicated that in pre-adipocytes TRAP 5a can interact with TRIP-1 intracellularly.

4.3 PAPER III

The aim in this paper was to determine if TRAP 5a can interact with the cell surface of pre-adipocytes and to investigate the routes through which TRAP 5a gets endocytosed by the cells. We also aimed to investigate TRAP 5a interaction with heparin and heparan sulfate and with the HPSG glypican-4 that has previously been suggested to interact with TRAP in bone ¹⁶⁴.

Firstly, pre-adipocytes were treated with TRAP 5a and TRAP 5b at 4°C to prevent endocytosis and using immunocytochemistry and electron microscopy it was shown that TRAP 5a but not TRAP 5b can bind to the plasma membrane of the cells in a dose-dependent manner as shown by FACS. The staining revealed a spot-like pattern that could indicate the formation of receptor clusters, a process known to be used by some receptors for activation ²⁰¹. When the same method was applied at 37°C where endocytosis is allowed, TRAP 5a was shown to be endocytosed in a dose-dependent manner. Interestingly, no TRAP 5a endocytosis was observed for differentiated adipocytes, indicating that the expression of a binding partner for TRAP 5a in pre-adipocytes is down-regulated during differentiation.

Using gel filtration chromatography and activity assays, it was shown that TRAP 5a forms complexes with heparin and heparan sulfate, however only heparin was able to inhibit TRAP 5a phosphatase activity. The binding of TRAP 5a to heparan sulfate is of particular interest since a lot of proteins that can have a mitogenic effect mediate their effect through binding to heparan sulfate in the ECM and cell surface, e.g. FGF2 binding to FGF2 receptor leads to dimerization that is regulated by binding to heparan sulfate^{202–204}.

Co-localization between Glypican-4 and TRAP 5a was addressed and it was shown that TRAP 5a was co-localized with glypican-4 on the cell surface membrane and on endocytic vesicles, indicating that TRAP 5a could potentially bind to the cell membrane using the HPSG glypican-4 as a co-receptor for signalling and/or endocytosis facilitation as observed for other glypicans and their ability to bind to growth factors and facilitate signalling from the cell surface^{205,206}.

As described above, TRAP 5a was shown to bind to pre-adipocyte cell surface and found in intracellular vesicles so next we aimed to investigate the endocytosis route of TRAP 5a. Proteins that act as growth factors or mitogens have been shown to follow to major pathways of endocytosis, i.e. the clathrin- or caveolae- mediated pathways that are both dependent on dynamin. Firstly we questioned if a dynamin inhibitor could prevent endocytosis and it was shown that TRAP 5a endocytosis is indeed dynamin- dependent. Subsequently, using clathrin- and caveolae- specific inhibitors it was shown that TRAP 5a endocytosis is caveolae-mediated, which is an endocytosis route used only by a few growth factors since most growth factors are endocytosed in a clathrin-dependent manner²⁰⁷. Interestingly, it has been shown that caveolae-mediated endocytosis is responsible for insulin signalling²⁰⁸ and that triacylglycerol synthesis takes place in the caveolae²⁰⁹, thus indicating that TRAP signalling could potentially act in a similar manner. However this could also depend on the concentration of TRAP 5a used, as in this study 100M were used- a quite excessive concentration and it has previously been reported that cell could use caveolae- dependent endocytosis in the presence of high concentrations of growth factors to target them to degradation, e.g. EGF that at low concentrations is endocytosed in a clathrin- mediated manner but at a high concentration instead by a clathrin- independent manner¹⁰.

TRAP 5a was shown to co-localize with the late endosomal marker Rab-7 and by electron microscopy to be also found in multivesicular bodies. However, it did not co-localize with the lysosomal marker LAMP-1, indicating that TRAP 5a either does not end up in the lysosomes or that it is degraded rapidly thus it is not detected, unlike what has been observed for TRAP in osteoblasts that co-localizes with LAMP-1¹⁵⁶, probably due to the different routes followed by the two cell types for TRAP processing since TRAP is expressed endogenously in osteoblasts^{149–153} but not in pre-adipocytes¹³⁸.

Finally, mesenchymal stem cells and pre-osteoblasts and fibroblasts (that are also of mesenchymal lineage) were studied for their ability to bind TRAP 5a and it was shown that pre-osteoblasts but not MSCs or fibroblasts had the ability to bind TRAP 5a. TRAP 5a binding by pre-osteoblasts is in line with previous findings that indicate that pre-osteoblasts are able to bind TRAP¹⁵³. Apart from pre-adipocytes TRAP 5a was shown to bind to pre-osteoblasts but not fibroblasts or mesenchymal stem cells.

In summary, this study aimed to identify if TRAP 5a binds to cell surface on pre-adipocytes and the route of endocytosis thereafter as well as to investigate if TRAP 5a colocalizes with the previous suggested binding partner glypican-4 and interaction with heparin and heparan sulfate. Results showed that TRAP 5a but not TRAP 5b binds to the cell surface of pre-adipocytes, gets endocytosed in a caveolae- dependent manner and ends-up in late endosomes and multivesicular bodies. Additionally it was shown that TRAP 5a forms complex with heparin and heparan sulfate and colocalizes with the HPSG glypican-4.

4.4 PAPER IV

The aim of paper IV was to investigate the effect of TRAP 5a exposure, mimicking a paracrine situation, on cell cycle, differentiation and migration of pre-adipocytes and to identify potential signalling pathways affected by TRAP 5a treatment in pre-adipocytes shortly after cell cycle entry.

Firstly, cells were incubated with serum- free media for cell cycle arrest and synchronization and then they were allowed to enter the cell cycle in the presence or absence of TRAP 5a. Results indicated an increased number of cells in the S phase after 12h (~50% more cells) and increased cyclin D1 expression.

Cyclin D1 expression is an important hallmark in cell cycle progression as it signifies the G1 to S transition²⁴ and is regulated by intracellular signals such as e.g. GSK3 β inactivation^{22,23} by activated Akt [40] or by Wnt- mediated signalling²⁶, thus next we aimed to screen for potential phosphorylation changes in proteins of the Akt signalling pathway using a protein phosphorylation array. Interestingly results showed a decreased phosphorylation in Akt and Erk sites responsible for its activation suggesting that perhaps the action of TRAP 5a on cell number regulation is not mediated via direct activation of Akt or that TRAP 5a leads to faster response by the cells and transient activating phosphorylations of Akt²⁰ to be already down-regulated already after 5 min.

On the other hand, GSK3 β showed reduced phosphorylation that leads to its inactivation and subsequent β -catenin stabilization that could account for effects on e.g. cyclin D1 levels and anti-apoptotic genes^{34,210,211}. Apart from Akt, GSK3 β regulation is mediated via canonical and non-canonical pathways of Wnt signalling²¹⁰ that is also involved in cell cycle entry and apoptosis reduction³⁴ and thus could be a potential signalling pathway affected by TRAP 5a. Stat3 activating phosphorylation was shown to be increased by TRAP 5a treatment and Stat3 has been shown to have an effect on the proliferation of pre-adipocytes, cyclin D1 levels and anti-apoptotic genes^{212,213}. After serum addition and cell cycle entry an up-regulation of phosphorylations leading to activation of pro-apoptotic proteins p53 and p38 in control cells was observed. Increased activity of pro-apoptotic proteins could be attributed to hypermitogenic stimulation²⁸. Interestingly, p53 and p38 exhibited reduced activating phosphorylation and Bad exhibited increased inhibiting phosphorylation after TRAP 5a treatment, indicating a potential effect of TRAP 5a in genes regulating cell cycle arrest and apoptosis, a mechanism that could also be involved in the cell number regulation as the 'break' in cell cycle progression. Activating phosphorylation of HSP27 that is involved in cell cycle arrest and is a downstream target of p38^{214,215} was also observed after TRAP 5a treatment, further suggesting a p38- and p53- mediated effect of TRAP 5a in cell cycle arrest. When p38 is activated cyclin D1 is inhibited thus leading to cell cycle arrest, while p53 is increased leading to p21 activation that results in cell cycle arrest or p16, p14/p19 activation resulting in premature senescence^{216–218}. In this system, cells have been serum- deprived and cell cycle-arrested, thus a reduced activation of p38 and p53 would likely indicate an increased evasion of the cell cycle arrest after serum stimulation.

Proteins p38, p53 and heat shock protein 27 (HSP27) have been shown to effect several biological events, such as migration^{215,219–222}, which can be also coupled to proliferation. This idea is contrary to previous beliefs that proliferation and migration are two separate events that cannot take place simultaneously^{223,224}. Since TRAP has previously been shown to have an effect on the migration of osteoclasts²²⁵ and migration and morphology of cancer cells¹⁸⁵, we investigated the effect of TRAP 5a on morphology, migration and protein expression of proteins related to migration adhesion and cell shape such as focal adhesion kinase (FAK), paxillin and actin. Morphological analysis and wound healing assay showed that TRAP 5a lead to less circular and more elongated cells and increased migration. Also, synchronized pre-

adipocytes were assessed for morphology after serum stimulation in the presence of TRAP 5a and it was shown that cells became larger at low concentrations of TRAP 5a but less circular and more elongated at high concentrations. Protein expression of FAK and paxillin was decreased after 8h of TRAP 5a treatment, whereas β -actin was decreased after 5h, indicating that apart from morphology, TRAP 5a could exert an effect on the adhesion pattern of the cells.

Finally, the effect of TRAP 5a in early differentiation of pre-adipocytes was tested and it was shown that TRAP 5a enhances lipid droplet acquisition already after 2 days of differentiation independently of stimulation by the PPAR γ activator rosiglitazone. The mechanism through which differentiation is affected by TRAP 5a remains unknown, however since pro-apoptotic proteins have been shown to have an effect on pre-adipocyte differentiation and differentiating pre-adipocytes exhibit increased resistance to apoptosis ²²⁶ it is likely that pre-adipocyte differentiation is stimulated by TRAP 5a via the same signalling pathway responsible for augmented G1 to S transition.

In summary, this study aimed to investigate the outcome of TRAP 5a treatment on pre-adipocytes and it was shown that TRAP 5a enhances the proliferation, migration and differentiation of pre-adipocytes. In a protein level, TRAP was shown to increase cyclin D1 levels and to change the phosphorylation pattern of GSK3 β and to decrease activating phosphorylation of pro-apoptotic proteins p53 and p38. Finally, TRAP 5a was shown to affect cell morphology and reduce the expression of β -actin, FAK and paxillin. These results suggest an overall effect in the cell cycle events by TRAP 5a that could possibly be mediated by a common pathways affecting GSK3 β activity.

4.5 GENERAL CONCLUSIONS AND DISCUSSION

In paper I it was shown that serum levels of TRAP 5a from obese females are lower than those from lean females. As discussed earlier, this was an interesting finding since TRAP levels in adipose tissue of obese women are elevated ¹⁹². This could indicate different processing mechanisms of TRAP 5a from adipose tissue in lean and obese states. TRAP 5a expression is increased but perhaps the deposition to the serum could be reduced and more TRAP 5a remains associated to the tissue where it can stimulate the proliferation and differentiation of pre-adipocytes. Also since TRAP 5a has been shown to be endocytosed by pre-adipocytes in paper II, TRAP 5a levels in adipose tissue could be maintained high by endocytosis in the tissue. It has been shown that the size of adipocytes and type of obesity (hyperplastic vs. hypertrophic obesity) in visceral and subcutaneous depots is important for the expression levels of cytokines as well as for the metabolic effects ¹⁰⁹, so the depot and type of obesity could also be related to the TRAP 5a levels in serum. Decreased levels of tissue in the serum of obese women seems also contradictory to a study performed in Chinese overweight males where it was shown that TRAP 5a is elevated in males with BMI 25-30 kg/m²¹⁹³. However, as discussed earlier the study of paper I differs from the previous study in the BMI grouping (study I was performed in a group of obese females with BMI ≥ 30 kg/m²) as well as sex. It is therefore possible that TRAP 5a levels increase in serum of overweight individuals but decrease in the obese state which is more severe.

TRAP 5a is reduced in the serum of obese females but increased in the adipose tissue. In paper II and paper III it was shown that TRAP 5a can bind to components of the adipose tissue such as nidogen-2 and heparan sulfate. As discussed earlier, proteins with mitogenic effects are often binding to ECM proteins to be delivered to the cells or to be stored until they are needed when ECM is degraded and they are released. Since TRAP 5a seems to have an effect on proliferation, differentiation and migration of pre-adipocytes, thus exhibiting growth factor properties, the biological implications of such interactions with components of the ECM could

serve the purpose of storage and later release by degradation of the ECM, or as to be delivered to the cells for binding on the cell surface.

Binding of TRAP 5a on the cell surface is accompanied by co-localization with the HPSG glypican-4, also suggesting that if the mitogenic effects of TRAP 5a are following a mechanism of action similar to growth factors, glypican-4 could act as a co-receptor for TRAP 5a binding. Glypican-4 has been shown to enhance the signalling of Wnt 5a and to get endocytosed together with Wnt receptor Frizzled-2²²⁷, which is consistent with the observation in paper III of co-localization of TRAP 5a with glypican-4 intracellularly and on cell surface, thus suggesting that the endocytosis and effect of TRAP 5a could potentially also be mediated in a similar manner as Wnt proteins. This hypothesis could further be strengthened by results from the signalling array in paper IV where reduced phosphorylation leading to inhibition of GSK3 β activity was observed after TRAP 5a exposure and as discussed previously, GSK3 β activity is mainly regulated by Akt and Wnt (canonical and non-canonical) signalling. Endocytosis of TRAP 5a occurs via the caveolae pathway and even though, generally caveolae pathway is a rare route for signalling process but rather for degradation it is responsible for the insulin endocytosis and triacylglycerol synthesis in adipocytes^{208,209} indicating an importance in adipocyte signalling. Binding of TRAP 5a was also observed for pre-osteoblasts, which is similar to what has been shown in previous studies for endocytosis of TRAP and effect on their differentiation^{151,153,187}.

TRAP 5a was also shown to affect p53, p38 and Bad phosphorylation pattern indicating that there might be a potential regulatory role of TRAP 5a in cell cycle arrest/ senescence signalling. These data suggest that TRAP 5a could affect both the 'gas' and 'break' mechanisms of cell cycle progression.

On cellular level TRAP 5a leads apart from increased proliferation to increased migration as well as differentiation which in normal tissue could be important for tissue repair. On the other hand in pathological situations, dysregulation of TRAP expression could potentially be responsible for cell transformation to a more proliferative and migrating phenotype as observed for cancer cells with increased TRAP levels.

4.6 REVISED WORKING HYPOTHESIS

In light of the data in this thesis and previous studies on TRAP 5a action in adipose tissue, a revised hypothesis of TRAP 5a action is described in figure 9. A possible mechanism of action would be that TRAP 5a is secreted by a subset of macrophages and interacts with components of the ECM such as HSPGs and nidogen-2 in order to be stored or delivered to pre-adipocytes in close vicinity. TRAP 5a on the cell surface of pre-adipocytes could bind to a yet unidentified receptor either directly or indirectly facilitated by glypican-4 and HSPGs. TRAP 5a endocytosis via the caveolae-mediated pathway and presence in multivesicular bodies and endosomes could trigger a signalling pathway mediated by changes in phosphorylation of GSK3 β , leading to various possible outcomes, such as increased cyclin D1 expression and G1 to S transition and migration and/or differentiation to adipocytes. These outcomes would be relevant for tissue homeostasis in a tissue repair/ wound healing situation. If dysregulated, eg overexpression, these outcomes could potentially lead to cell transformation.




Figure 9. Revised working hypothesis. TRAP 5a secreted from macrophages in adipose tissue binds to the ECM probably by interaction with nidogen-2 and heparin or heparan sulfate. TRAP 5a binds to the surface of pre-adipocytes in proximity to the HSPG glypican-4 and is endocytosed via the caveolae pathway. Intracellularly it is found in early endosomes and multivesicular bodies but is not co-localized with LAMP-1. TRAP 5a enhances the proliferation of pre-adipocytes and increases cyclin D1 levels, increases differentiation (assessed by lipid droplet acquisition) and causes morphological changes and migration enhancement and decrease in FAK, paxillin and actin.

With regard to this hypothesis the following question would be relevant: How can such an action from a macrophage- secreted molecule contribute to tissue homeostasis? TRAP 5a mutations in humans have been shown to lead to bone dysplasia and autoimmune disorders, however there have not been any published observations of complete lack of adipose tissue formation^{174,177}, thus indicating that TRAP 5a is not critical for embryonic adipose tissue development. An alternative explanation could be that TRAP 5a contributes to tissue homeostasis in mature adipose tissue in cases where tissue repair or regeneration is needed. As previously mentioned, adipose tissue exhibits a stable number of adipocyte cells and an annual turnover of 10%^{55,56}, however in obesity there is an increased rate of adipocyte cell death and increased number of macrophages⁹⁸, suggesting an increased need for tissue ‘repair’. Thus, one explanation of the biological relevance of TRAP 5a action would be as a part of the tissue repair (a wound healing-like action) of adipose tissue in obesity, triggered by macrophages. Similar mechanism have been previously shown for macrophages in other types of tissue such as skin where M2 anti-inflammatory macrophages express growth factors and stimulate cell differentiation and tissue repair in injury sites⁹⁶ and TRAP 5a could potentially be a part of the tissue repair mechanism in adipose tissue. However, several aspects of this hypothesis need to be further investigated.

4.7 FUTURE PERSPECTIVES AND CLINICAL SIGNIFICANCE

The revised hypothesis on TRAP 5a action presented above, would indicate several areas of research regarding TRAP 5a action.

Up to today, the subset of macrophages that express TRAP remains unknown and as previously described, macrophages can adopt a pro-inflammatory (M1) or anti-inflammatory (M2) phenotype as well as intermediate stages. Identification of the phenotype that expresses TRAP 5a would further contribute to the understanding of the role of TRAP 5a in adipose tissue. Additionally stimuli of TRAP expression from macrophages e.g. hypoxia, free fatty acids etc. that participate in macrophage recruitment and activation in adipose tissue remain to be investigated. Moreover the finding that TRAP 5a serum levels are reduced in the obese state, could be further studied by correlating TRAP 5a levels to type of obesity (e.g. hyperplastic or hypertrophic), adipocyte size and number. The newly developed TRAP 5a ELISA could be used to further investigate the use of this specific isoform as a marker for disease.

The difference between the functions two isoforms is of particular interest as it represents a situation where proteolytical cleavage switches the function from a protein with mitogenic/growth factor-like effect (TRAP 5a) to a phosphatase (TRAP 5b). Thus the conditions under which this transition occurs would be of interest in future studies.

Regarding TRAP 5a action in pre-adipocytes, one important future aspect would be the identification of the signalling binding partner of TRAP 5a on the pre-adipocyte cell surface. Identification of this binding partner would allow for further targeting of TRAP 5a action and future use for screening in other tissues for similar action.

Additionally, the exact signalling pathway of TRAP 5a in pre-adipocytes remains to be identified. A future study could involve the testing of hypothesis of wnt signalling- mediated TRAP action by the use of wnt signalling receptor inhibitors.

Finally the hypothesis of TRAP 5a acting as a part of tissue repair mechanism could be further tested in other tissues of mesenchymal origin. Studies suggest similar effects in pre-osteoblasts thus the effect of TRAP 5a on chondrocytes could also be of interest.

In summary, further elucidation of TRAP 5a mechanism action by identification of the binding partner and signalling pathway could contribute to the development of a drug target for hindering increased proliferation, migration and differentiation of pre-adipocytes and thus adipose tissue development. Moreover, TRAP 5a mechanism of action could be useful for the development of methods to promote adipose tissue regeneration in pathological situations such as cachexia where absence of adipose tissue can be life threatening. Additionally, it is of importance to investigate if the mechanism applies to other normal tissue types such as bone and cartilage or in cancer tissues where TRAP has also been shown to play a role.

5 ACKNOWLEDGEMENTS

The work presented in this thesis was financially supported by grants from the **Swedish Research Council** and the Research Funds of **Karolinska Institutet**.

Since I started my work at the division of Pathology, I have met many wonderful people that have helped me improve as a researcher but also personally. I would like to express my gratitude to the following people:

First of all, I would like to thank my main supervisor **Göran Andersson** and my co-supervisor **Pernilla Lång**. **Göran** thank you for giving me the opportunity to be your PhD student, for continuously inspiring me with your enthusiasm for science and bright ideas and for encouraging me to develop my own work. I have enjoyed all the interesting and fun scientific discussions we had during the whole time I have been in your group. **Pernilla**, thank you for helping me develop as a researcher, for always supporting me when I got discouraged and for motivating me when I needed it the most. I admire your enthusiasm for science and I have learned a lot from you.

Big thank you to **Maria Norgård** and **Barbro Ek-Rylander** for all the good meetings I had with you all these years and for your help and feedback. I would also like to thank **Annika Nordvall** for her help and feedback from the beginning of this thesis, **Erik Karlström** and **Serhan Zenger** for helping me on my first days at the division of Pathology and especially **Carolina Wejheden** for not only helping me every day in the lab but also for being a good friend.

I have also had the opportunity to help with the supervision of students during my thesis. A big thank you **Sara Witte**, **Tania Brandstötter**, **Christina Redmers**, **Hong Anh Mai**, **Heike Becker** and **Eva van der Heijden** for bringing fresh ideas, great skills and helping me with the completion for this thesis.

I would also like to thank **Peter Arner** and **Staffan Paulie** for a very good collaboration. Special thank you to **Birgitta Sander** and **Xiao Wang** for a collaboration that has taught me a lot. I would also like to thank **Åsa-Lena Dackland** for her help with FACS, **Birger Christensson** for the help with the confocal microscope, **Kjell Hultenby** for the help with the electron microscopy and **Gareth Morgan** for his help during teaching and sharing his knowledge. I would also like to thank **Lennart Eriksson**, **Mikael Björnstedt**, **Aristi Fernandes**, **Katalin Dobra**, **Anders Hjerpe**, **Stephen Strom**, for their feedback and discussions during PhD seminars, **Andrej Weintraub** and **Marita Ward** for their help in the PhD process and **Arja Kramsu** for her help during teaching and **Eva Hjertquist Duräng**, **Annika Käller** and **Helena Andersson** for their valuable administrative help

Time flies when you are having fun and the years while performing this work went by very fast thanks to the great working environment. I would like to thank **Eric Olm**, **Markus Selenius**, and **Stefan Almestrand** for the fun times at the division and retreats, **Marita Wallenberg**, **Rim Jawad**, **Sougat Misra**, **Massoud Vosough**, **Kristina Kannisto**, **Roberto**

Gramignoli, Carl-Olof Hilledal, Raoul Kuiper, Michael Krumpel, Dilruba Ahmed, Sara Arroyo Muhr, Shuvojit Moulik, Rita Hutyra-Gram Ötvös I would like to thank you for all the fun during lunch, retreats and fika! I would also like to thank **Ghazal Heidari-Hamedani** and **Tünde Szatmári** for the great talks at work and the time in Greece, **Magali Merrien** and **Teresa Seestern** for their positive attitude and friendship, **Laia Mira Pasqual** for her great attitude, sharp ideas and for the help with the ELISAs, **Martin Lord** for motivation with running and funny comments, **Annelie Mollbrink** for making the work environment great, **Hannes Olauson** for all the pleasant talks in lab and at lunch, **Vicky Chatzakos** for her positive attitude and for being a great friend, **Lisa Arodin** for sharing all the PhD experience, for friendship since our first PhD student days and all the great time outside work! I would like to thank **Adam Szulkin** for helping me with teaching, for the valuable advice and for all the laughs. To **Anja Reithmeier** a big thank you for having someone to share the lab troubles, the same taste in music and the fun times outside work. I would like to especially thank **Agata Wasik** for always being supportive, helpful and for her great working attitude and friendship and **Filip Mundt** for all the great talks, his help during the writing of this thesis and for being a good friend.

Apart from the division of Pathology I would like to thank **Eleftheria Palkopoulou** and **Sewit Mogos** for their friendship all these years, **Barry Whelan** for reminding me to take breaks during the writing of this thesis **Athina Manolia-Bär, Sebastian Bär and Dimitris Bär** for always supporting me and **Yannis Bistinas** for sharing the PhD concerns.

The biggest thank you goes to my brother **Panagiotis Patlakas** for always supporting me and encouraging me to reach my goals and being always there to help me and my parents **Triantafyllos** and **Despoina** for their constant support and help, this thesis would not have been possible without you. Το μεγαλύτερο ευχαριστώ το οφείλω στους γονείς μου για την υποστήριξή τους και την βοήθεια τους όποτε τη χρειάζομαι.

6 REFERENCES

1. Robert, L. The Extracellular Matrix. *FEBS Lett.* **268**, 386–393 (1990).
2. Cristancho, A. G. & Lazar, M. a. Forming functional fat: a growing understanding of adipocyte differentiation. *Nat. Rev. Mol. Cell Biol.* **12**, 722–734 (2011).
3. Gurevich, V. V *et al.* Arrestin interactions with G protein-coupled receptors. Direct binding studies of wild type and mutant arrestins with rhodopsin, beta 2-adrenergic, and m2 muscarinic cholinergic receptors. *J. Biol. Chem.* **270**, 720–731 (1995).
4. Orlichenko, L., Huang, B., Krueger, E. & McNiven, M. a. Epithelial growth factor-induced phosphorylation of caveolin 1 at tyrosine 14 stimulates caveolae formation in epithelial cells. *J. Biol. Chem.* **281**, 4570–4579 (2006).
5. Murata, M. *et al.* VIP21/caveolin is a cholesterol-binding protein. *Proc. Natl. Acad. Sci. U. S. A.* **92**, 10339–43 (1995).
6. Damke, H., Baba, T., Warnock, D. E. & Schmid, S. L. Induction of mutant dynamin specifically blocks endocytic coated vesicle formation. *J. Cell Biol.* **127**, 915–934 (1994).
7. Hill, E., van Der Kaay, J., Downes, C. P. & Smythe, E. The role of dynamin and its binding partners in coated pit invagination and scission. *J. Cell Biol.* **152**, 309–23 (2001).
8. Platta, H. W. & Stenmark, H. Endocytosis and signaling. *Curr. Opin. Cell Biol.* **23**, 393–403 (2011).
9. Poteryaev, D., Datta, S., Ackema, K., Zerial, M. & Spang, A. Identification of the switch in early-to-late endosome transition. *Cell* **141**, 497–508 (2010).
10. Sigismund, S. *et al.* Clathrin-independent endocytosis of ubiquitinated cargos. *Proc. Natl. Acad. Sci. U. S. A.* **102**, 2760–2765 (2005).
11. Yamamoto, H., Sakane, H., Yamamoto, H., Michiue, T. & Kikuchi, A. Wnt3a and Dkk1 Regulate Distinct Internalization Pathways of LRP6 to Tune the Activation of β -Catenin Signaling. *Dev. Cell* **15**, 37–48 (2008).
12. Palamidessi, A. *et al.* Endocytic Trafficking of Rac Is Required for the Spatial Restriction of Signaling in Cell Migration. *Cell* **134**, 135–147 (2008).
13. Hayes, S., Chawla, A. & Corvera, S. TGF β receptor internalization into EEA1-enriched early endosomes: Role in signaling to Smad2. *J. Cell Biol.* **158**, 1239–1249 (2002).
14. Tsukazaki, T., Chiang, T. a., Davison, A. F., Attisano, L. & Wrana, J. L. SARA, a FYVE domain protein that recruits Smad2 to the TGF β receptor. *Cell* **95**, 779–791 (1998).

15. Dobrowolski, R. & De Robertis, E. M. Endocytic control of growth factor signalling: multivesicular bodies as signalling organelles. *Nat. Rev. Mol. Cell Biol.* **13**, 53–60 (2012).
16. Mellman, I. & Warren, G. The road taken: past and future foundations of membrane traffic. *Cell* **100**, 99–112 (2000).
17. Nickel, W. & Rabouille, C. Mechanisms of regulated unconventional protein secretion. *Nat. Rev. Mol. Cell Biol.* **10**, 148–155 (2009).
18. Wegehangel, S., Zehe, C. & Nickel, W. Rerouting of fibroblast growth factor 2 to the classical secretory pathway results in post-translational modifications that block binding to heparan sulfate proteoglycans. *FEBS Lett.* **582**, 2387–2392 (2008).
19. Pardee, a B. G1 events and regulation of cell proliferation. *Science* **246**, 603–608 (1989).
20. Manning, B. D. & Cantley, L. C. AKT/PKB signaling: navigating downstream. *Cell* **129**, 1261–74 (2007).
21. Papkoff, J. & Aikawa, M. WNT-1 and HGF regulate GSK3 beta activity and beta-catenin signaling in mammary epithelial cells. *Biochem. Biophys. Res. Commun.* **247**, 851–8 (1998).
22. Alao, J. P. The regulation of cyclin D1 degradation: roles in cancer development and the potential for therapeutic invention. *Mol. Cancer* **6**, 24 (2007).
23. Shtutman, M. *et al.* The cyclin D1 gene is a target of the χ -catenin TEF-1 pathway. *Pro. Natl. Acad. Sci. USA* **96**, 5522–5527 (1999).
24. Liang, J. & Slingerland, J. M. Multiple Roles of the PI3K / PKB (Akt) Pathway in Cell Cycle Progression *Cell Cycle* **2:4**, 339-345 (2003).
25. Cross, D., Alessi, D., Cohen, P., Andjelkovic, M. & Hemmings, B. Inhibition of glycogen synthase kinase-3 by insulin mediated by protein kinase B. *Nature* **21-28**, 785-9 (1995).
26. Papkoff, J. & Aikawa, M. WNT-1 and HGF Regulate GSK3 b Activity and b - Catenin Signaling in Mammary Epithelial Cells. *Biochem Biophys Res Commun.* **247**, 851–858 (1998).
27. Gad, A., Thullberg, M., Dannenberg, J.-H., te Riele, H. & Strömblad, S. Retinoblastoma susceptibility gene product (pRb) and p107 functionally separate the requirements for serum and anchorage in the cell cycle G1-phase. *J. Biol. Chem.* **279**, 13640–4 (2004).
28. Blagosklonny MV ., Cell Senescence : Hypertrophic Arrest Beyond the Restriction Point. *J Cell Physiol.* , **597**, 592–597 (2006).
29. Ohtani, N. & Hara, E. Roles and mechanisms of cellular senescence in regulation of tissue homeostasis. *Cancer Sci.* **104**, 525–30 (2013).

30. Muñoz-Espín, D. & Serrano, M. Cellular senescence: from physiology to pathology. *Nat. Rev. Mol. Cell Biol.* **15**, 482–96 (2014).
31. Campisi, J. & d’Adda di Fagagna, F. Cellular senescence: when bad things happen to good cells. *Nat. Rev. Mol. Cell Biol.* **8**, 729–740 (2007).
32. Wada, T. & Penninger, J. M. Mitogen-activated protein kinases in apoptosis regulation. *Oncogene* **23**, 2838–49 (2004).
33. Hipfner, D. R. & Cohen, S. M. Connecting proliferation and apoptosis in development and disease. *Nat. Rev. Mol. Cell Biol.* **5**, 805–815 (2004).
34. You, Z. *et al.* Wnt signaling promotes oncogenic transformation by inhibiting c-Myc-induced apoptosis. *J. Cell Biol.* **157**, 429–40 (2002).
35. Imhof, B. a & Aurrand-Lions, M. Adhesion mechanisms regulating the migration of monocytes. *Nat. Rev. Immunol.* **4**, 432–444 (2004).
36. Millard, M., Odde, S. & Neamati, N. Integrin targeted therapeutics. *Theranostics* **1**, 154–188 (2011).
37. Legate, K. R., Wickström, S. a, Fässler, R., Fa, R. & Wickstro, S. a. Genetic and cell biological analysis of integrin outside-in signaling Genetic and cell biological analysis of integrin outside-in signaling. *Genes Dev.* **23**, 397–418 (2009).
38. Smart, N. & Riley, P. R. The stem cell movement. *Circ. Res.* **102**, 1155–1168 (2008).
39. Takada, I., Kouzmenko, A. P. & Kato, S. Wnt and PPARgamma signaling in osteoblastogenesis and adipogenesis. *Nat. Rev. Rheumatol.* **5**, 442–7 (2009).
40. Huang, H. *et al.* BMP signaling pathway is required for commitment of C3H10T1/2 pluripotent stem cells to the adipocyte lineage. *Proc. Natl. Acad. Sci. U. S. A.* **106**, 12670–5 (2009).
41. Brizzi, M. F., Tarone, G. & Defilippi, P. Extracellular matrix, integrins, and growth factors as tailors of the stem cell niche. *Curr. Opin. Cell Biol.* **24**, 645–51 (2012).
42. He, X., Semenov, M., Tamai, K. & Zeng, X. LDL receptor-related proteins 5 and 6 in Wnt/beta-catenin signaling: arrows point the way. *Development* **131**, 1663–1677 (2004).
43. Bennett, C. N. *et al.* Regulation of Wnt signaling during adipogenesis. *J. Biol. Chem.* **277**, 30998–1004 (2002).
44. Gregoire, F. M., Smas, C. M. & Sul, H. S. Understanding adipocyte differentiation. *Physiol. Rev.* **78**, 783–809 (1998).
45. Miettinen, S., Sarkanen, J. R. & Ashammakhi, N. Adipose Tissue and Adipocyte Differentiation: Molecular and Cellular Aspects and Tissue Engineering Applications. *Top. Tissue Eng.* **4**, 1–26 (2008).

46. Mohamed-Ali, V., Pinkney, J. H. & Coppack, S. W. Adipose tissue as an endocrine and paracrine organ. *Int. J. Obes. Relat. Metab. Disord.* **22**, 1145–1158 (1998).
47. Flier, J. S. What 's in a Name ? In Search of Leptin 's Physiologic Role. *Endocrinol. Metab.* **83**, 1407–1413 (1998).
48. Kim, S. & Moustaid-moussa, N. Symposium : Adipocyte Function , Differentiation and Metabolism Secretary , Endocrine and Autocrine / Paracrine Function of the Adipocyte 1. *J. Nutr.* 3110–3115 (2000).
49. Greenberg, A. S. *et al.* Perilipin, a major hormonally regulated adipocyte-specific phosphoprotein associated with the periphery of lipid storage droplets. *J. Biol. Chem.* **266**, 11341–11346 (1991).
50. Ono, M., Aratani, Y., Kitagawa, I. & Kitagawa, Y. Ascorbic acid phosphate stimulates type IV collagen synthesis and accelerates adipose conversion of 3T3-L1 cells. *Exp. Cell Res.* **187**, 309–314 (1990).
51. Aratani, Y. & Kitagawa, Y. Enhanced synthesis and secretion of type IV collagen and entactin during adipose conversion of 3T3-L1 cells and production of unorthodox laminin complex. *J. Biol. Chem.* **263**, 16163–9 (1988).
52. Sillat, T. *et al.* Basement membrane collagen type IV expression by human mesenchymal stem cells during adipogenic differentiation. *J. Cell. Mol. Med.* **16**, 1485–95 (2012).
53. Molina, H. *et al.* Temporal profiling of the adipocyte proteome during differentiation using a five-plex SILAC based strategy *J Proteome Res.* **8**, 48–58 (2010).
54. Poissonnet, C. M., Burdi, a R. & Bookstein, F. L. Growth and development of human adipose tissue during early gestation. *Early Hum. Dev.* **8**, 1–11 (1983).
55. Arner, E. *et al.* Adipocyte Turnover : Relevance to Human Adipose Tissue Morphology. *Production* **59**, 105–109 (2010).
56. Spalding, K. L. *et al.* Dynamics of fat cell turnover in humans. *Nature* **453**, 783–7 (2008).
57. Wernstedt Asterholm, I. *et al.* Adipocyte inflammation is essential for healthy adipose tissue expansion and remodeling. *Cell Metab.* **20**, 103–118 (2014).
58. Friedenstein, a J., Piatetzky-Shapiro, I. I. & Petrakova, K. V. Osteogenesis in transplants of bone marrow cells. *J. Embryol. Exp. Morphol.* **16**, 381–90 (1966).
59. Pittenger, M. F. Multilineage Potential of Adult Human Mesenchymal Stem Cells. *Science (80-.).* **284**, 143–147 (1999).
60. Chamberlain, G., Fox, J., Ashton, B. & Middleton, J. Concise review: mesenchymal stem cells: their phenotype, differentiation capacity, immunological features, and potential for homing. *Stem Cells* **25**, 2739–49 (2007).

61. Takada, I., Suzawa, M., Matsumoto, K. & Kato, S. Suppression of PPAR transactivation switches cell fate of bone marrow stem cells from adipocytes into osteoblasts. *Ann. N. Y. Acad. Sci.* **1116**, 182–95 (2007).
62. Takada, I., Kouzmenko, A. P. & Kato, S. Molecular switching of osteoblastogenesis versus adipogenesis: implications for targeted therapies. *Expert Opin. Ther. Targets* **13**, 593–603 (2009).
63. Li, C. *et al.* MicroRNA-188 regulates age-related switch between osteoblast and adipocyte differentiation. *J Clin Invest.* **57127**, 1–14
64. Schiller, P. C., D'Ippolito, G., Brambilla, R., Roos, B. a. & Howard, G. a. Inhibition of Gap-Junctional Communication Induces the Trans-differentiation of Osteoblasts to an Adipocytic Phenotype in Vitro. *J. Biol. Chem.* **276**, 14133–14138 (2001).
65. Ono, N., Ono, W., Nagasawa, T. & Kronenberg, H. M. A subset of chondrogenic cells provides early mesenchymal progenitors in growing bones. *Nat. Cell Biol.* **16**, 1157–1167 (2014).
66. Zhou, X. *et al.* Chondrocytes Transdifferentiate into Osteoblasts in Endochondral Bone during Development, Postnatal Growth and Fracture Healing in Mice. *PLoS Genet.* **10**, e1004820 (2014).
67. Green, H. & Meuth, M. An established pre-adipose cell line and its differentiation in culture. *Cell* **3**, 127–33 (1974).
68. Tang, W. *et al.* White fat progenitor cells reside in the adipose vasculature. *Science* **322**, 583–586 (2008).
69. Hudak, C. S. & Sul, H. S. Pref-1, a gatekeeper of adipogenesis. *Front. Endocrinol. (Lausanne)*. **4**, 79 (2013).
70. Prins, J. B. & O'Rahilly, S. Regulation of adipose cell number in man. *Clin. Sci. (Lond)*. **92**, 3–11 (1997).
71. Cryer, a, Williams, S. E. & Cryer, J. Dietary and other factors involved in the proliferation, determination and differentiation of adipocyte precursor cells. *Proc. Nutr. Soc.* **51**, 379–385 (1992).
72. Cornelius, P., Macdougald, O. A. & Lane, M. D. Regulation of adipocyte development. *Annu Rev Nutr.* **14**, 99–129 (1994).
73. Wright, J. T. & Hausman, G. J. Insulinlike growth factor-1 (IGF-1)-induced stimulation of porcine preadipocyte replication. *In Vitro Cell. Dev. Biol. Anim.* **31**, 404–8 (1995).
74. Hausman, D. B., Digirolamo, M., Bartness, T. J., Hausman, G. J. & Martin, R. J. The biology of white adipocyte proliferation. *Obes Rev*, **2**, 239–254 (2001).
75. Nikolai, a T., Wilde, M., Nakanishi, M., Smith, J. R. & Darlington, G. J. CCAAT / enhancer-binding protein (C / EBP) inhibits cell proliferation protein. **21**, 804–815 (1996).

76. Umek, R. M., Friedman, a D. & McKnight, S. L. CCAAT-enhancer binding protein: a component of a differentiation switch. *Science* **251**, 288–292 (1991).
77. Altioek, S., Xu, M. & Spiegelman, B. M. PPAR γ induces cell cycle withdrawal: Inhibition of E2f/DP DNA-binding activity via down-regulation of PP2A. *Genes Dev.* **11**, 1987–1998 (1997).
78. Richon, V. M., Lyle, R. E. & McGehee, R. E. Regulation and expression of retinoblastoma proteins p107 and p130 during 3T3-L1 adipocyte differentiation. *J. Biol. Chem.* **272**, 10117–10124 (1997).
79. Reichert, M. & Eick, D. Analysis of cell cycle arrest in adipocyte differentiation. *Oncogene*. **18**, 459-66 (1999).
80. Spiegelman, B. M. & Farmer, S. R. Decreases in tubulin and actin gene expression prior to morphological differentiation of 3T3 adipocytes. *Cell* **29**, 53–60 (1982).
81. Weiner, F. R., Shah, a, Smith, P. J., Rubin, C. S. & Zern, M. a. Regulation of collagen gene expression in 3T3-L1 cells. Effects of adipocyte differentiation and tumor necrosis factor alpha. *Biochemistry* **28**, 4094–9 (1989).
82. Peng, X.-D. *et al.* Dwarfism, impaired skin development, skeletal muscle atrophy, delayed bone development, and impeded adipogenesis in mice lacking Akt1 and Akt2. *Genes Dev.* **17**, 1352–65 (2003).
83. Rochford, J. J. Molecular mechanisms controlling human adipose tissue development: insights from monogenic lipodystrophies. *Expert Rev. Mol. Med.* **12**, e24 (2010).
84. Baudry, A., Yang, Z.-Z. & Hemmings, B. a. PKB α is required for adipose differentiation of mouse embryonic fibroblasts. *J. Cell Sci.* **119**, 889–97 (2006).
85. Rosen, E. D. & MacDougald, O. a. Adipocyte differentiation from the inside out. *Nat. Rev. Mol. Cell Biol.* **7**, 885–96 (2006).
86. Ibrahimi, a., Bonino, F., Bardon, S., Ailhaud, G. & Dani, C. Essential role of collagens for terminal differentiation of preadipocytes. *Biochem. Biophys. Res. Commun.* **187**, 1314–1322 (1992).
87. Galic, S., Oakhill, J. S. & Steinberg, G. R. Molecular and Cellular Endocrinology Adipose tissue as an endocrine organ. *Mol. Cell. Endocrinol.* **316**, 129–139 (2010).
88. Morris, D. L., Singer, K. & Lumeng, C. N. Adipose tissue macrophages: phenotypic plasticity and diversity in lean and obese states. *Curr. Opin. Clin. Nutr. Metab. Care* **14**, 341–6 (2011).
89. Patel, P. S., Buras, E. D. & Balasubramanyam, A. The role of the immune system in obesity and insulin resistance. *J. Obes.* **2013**, 616193 (2013).
90. Curat, C. A. *et al.* From Blood Monocytes to Adipose Tissue–Resident Macrophages. *Obes. Res.* **53**, (2004).

91. Jenkins, S. J. *et al.* Local macrophage proliferation, rather than recruitment from the blood, is a signature of TH2 inflammation. *Science* **332**, 1284–1288 (2011).
92. Lumeng, C. N., Bodzin, J. L. & Saltiel, A. R. Obesity induces a phenotypic switch in adipose tissue macrophage polarization. *J Clin Invest.* **117**, 175–84 (2007).
93. Lumeng, C. N., Delproposto, J. B., Westcott, D. J. & Saltiel, A. R. Phenotypic Switching of Adipose Tissue Macrophages With Obesity Is Generated by Spatiotemporal Differences in Macrophage Subtypes. *Diabetes* **57**, 3239–3246 (2008).
94. Suganami, T., Nishida, J. & Ogawa, Y. A paracrine loop between adipocytes and macrophages aggravates inflammatory changes: role of free fatty acids and tumor necrosis factor alpha. *Arterioscler. Thromb. Vasc. Biol.* **25**, 2062–8 (2005).
95. Odegaard, J. I. *et al.* Macrophage-specific PPARgamma controls alternative activation and improves insulin resistance. *Nature* **447**, 1116–1120 (2007).
96. Murray, P. J. & Wynn, T. a. Protective and pathogenic functions of macrophage subsets. *Nat. Rev. Immunol.* **11**, 723–737 (2011).
97. Suganami, T. & Ogawa, Y. Adipose tissue macrophages: their role in adipose tissue remodeling. *J. Leukoc. Biol.* **88**, 33–9 (2010).
98. Klimcakova, E. *et al.* Macrophage gene expression is related to obesity and the metabolic syndrome in human subcutaneous fat as well as in visceral fat. *Diabetologia* **54**, 876–87 (2011).
99. Romacho, T., Elsen, M., Röhrborn, D. & Eckel, J. Adipose tissue and its role in organ crosstalk. *Acta Physiol.* **210**, 733–753 (2014).
100. Weisberg, S. P. *et al.* Obesity is associated with macrophage accumulation. *J. Clin. Invest.* **112**, 1796–808 (2003).
101. Strissel, K. J. *et al.* Adipocyte death, adipose tissue remodeling, and obesity complications. *Diabetes* **56**, 2910–8 (2007).
102. Kabon, B. *et al.* Obesity decreases perioperative tissue oxygenation. *Anesthesiology* **100**, 274–80 (2004).
103. Dandona, P., Aljada, A. & Bandyopadhyay, A. Inflammation: the link between insulin resistance, obesity and diabetes. *Trends Immunol.* **25**, 4–7 (2004).
104. Hotamisligil, G. S. Inflammation and metabolic disorders. *Nature* **444**, 860–7 (2006).
105. Kamei, N. *et al.* Overexpression of monocyte chemoattractant protein-1 in adipose tissues causes macrophage recruitment and insulin resistance. *J. Biol. Chem.* **281**, 26602–14 (2006).

106. Kanda, H. *et al.* MCP-1 contributes to macrophage infiltration into adipose tissue, insulin resistance, and hepatic steatosis in obesity. *J. Clin. Invest.* **116**, 1494–505 (2006).
107. Watanabe, Y., Nagai, Y. & Takatsu, K. Activation and regulation of the pattern recognition receptors in obesity-induced adipose tissue inflammation and insulin resistance. *Nutrients* **5**, 3757–3778 (2013).
108. Hirsch, J. & Batchelor, B. Adipose tissue cellularity in human obesity. *Clin. Endocrinol. Metab.* **5**, 299–311 (1976).
109. Hoffstedt, J. *et al.* Regional impact of adipose tissue morphology on the metabolic profile in morbid obesity. *Diabetologia* **53**, 2496–2503 (2010).
110. Faust, I. M., Johnson, P. R., Stern, J. S. & Hirsch, J. Diet-induced adipocyte number increase in adult rats: a new model of obesity. *Am. J. Physiol.* **235**, E279–86 (1978).
111. Marques, B. G., Hausman, D. B. & Martin, R. J. Association of fat cell size and paracrine growth factors in development of hyperplastic obesity. *Am. J. Physiol.* **275**, R1898–908 (1998).
112. Ketcham, C. M., Roberts, R. M., Simmen, R. C. & Nick, H. S. Molecular cloning of the type 5, iron-containing, tartrate-resistant acid phosphatase from human placenta. *J. Biol. Chem.* **264**, 557–63 (1989).
113. Vincent JB, Averill BA. An enzyme with a double identity: purple acid phosphatase and tartrate-resistant acid phosphatase. *FASEB J.* **4**, 3009-14 (1990) .
114. Guddat, L. W. *et al.* Crystal structure of mammalian purple acid phosphatase. *Structure* **7**, 757–767 (1999).
115. Uppenberg, J., Lindqvist, F. & Svensson, C. Crystal Structure of a Mammalian Purple Acid Phosphatase. *J Mol Biol.* **290**, 201-11 (1999).
116. Sundararajan, T. A. & Sarma, P. S. Purification and properties of phosphoprotein phosphatase from ox spleen. *Biochem. J.* **56**, 125–30 (1954).
117. Li CY, Yam LT, Lam KW. Acid phosphatase isoenzyme in human leukocytes in normal and pathologic conditions *J Histochem Cytochem* **18**, 473-81(1970)
118. Li, C. Y., Chuda, R. A., Lam, W. K. & Yam, L. T. Acid phosphatases in human plasma. *J. Lab. Clin. Med.* **82**, 446–60 (1973).
119. Yam LT, Li CY, Lam KW. Tartrate-resistant acid phosphatase isoenzyme in the reticulum cells of leukemic reticuloendotheliosis. *N Engl J Med.* **284**, 357-60 (1971).
120. Halleen, J. M. *et al.* Tartrate-resistant acid phosphatase 5b: a novel serum marker of bone resorption. *J. Bone Miner. Res.* **15**, 1337–45 (2000).
121. Halleen, J. M. *et al.* Serum tartrate-resistant acid phosphatase 5b is a specific and sensitive marker of bone resorption. *Clin. Chem.* **47**, 597–600 (2001).

122. Halleen, J. M. *et al.* Serum tartrate-resistant acid phosphatase 5b, but not 5a, correlates with other markers of bone turnover and bone mineral density. *Calcif. Tissue Int.* **71**, 20–5 (2002).
123. Lord, D. K. *et al.* Type 5 acid phosphatase. Sequence, expression and chromosomal localization of a differentiation-associated protein of the human macrophage. *Eur. J. Biochem.* **189**, 287–93 (1990).
124. Leach RJ, Reus BE, Hundley JE, Johnson-Pais TL, Windle JJ. Confirmation of the assignment of the human tartrate-resistant acid phosphatase gene (ACP5) to chromosome 19. *Genomics.* **19**, 180-1.(1994)
125. Grimes R, Reddy SV, Leach RJ, Scarcez T, Roodman GD, Sakaguchi AY, Lalley PA, Windle JJ. Assignment of the mouse tartrate-resistant acid phosphatase gene (Acp5) to chromosome 9. *Genomics.* **15**, 421-2 (1993)
126. Cassady, A. I., King, A. G., Cross, N. C. & Hume, D. A. Isolation and characterization of the genes encoding mouse and human type-5 acid phosphatase. *Gene* **130**, 201–7 (1993).
127. Fleckenstein, E. & Drexler, H. G. Tartrate-resistant acid phosphatase: gene structure and function. *Leukemia* **11**, 10–3 (1997).
128. Walsh, N. C. *et al.* Multiple tissue-specific promoters control expression of the murine tartrate-resistant acid phosphatase gene. *Gene* **307**, 111–23 (2003).
129. Pan, W. *et al.* Analysis of distinct tartrate-resistant acid phosphatase promoter regions in transgenic mice. *J. Biol. Chem.* **280**, 4888–93 (2005).
130. Reddy, S. V *et al.* Characterization of the mouse tartrate-resistant acid phosphatase (TRAP) gene promoter. *J. Bone Miner. Res.* **10**, 601–6 (1995).
131. Lam, W. K. W., Eastlund, D. T., Li, C. & Yam, L. T. Biochemical Properties of Tartrate-Resistant Acid Phosphatase in Serum of Adults and Children. *Clin Chem.* **24**, 1105–1108 (1978).
132. Lam, K. W., Lee, P., Li, C. Y. & Yam, L. T. Immunological and biochemical evidence for identity of tartrate-resistant isoenzymes of acid phosphatases from human serum and tissues. *Clin. Chem.* **26**, 420–422 (1980).
133. Ljusberg, J. *et al.* Proteolytic excision of a repressive loop domain in tartrate-resistant acid phosphatase by cathepsin K in osteoclasts. *J. Biol. Chem.* **280**, 28370–81 (2005).
134. Ljusberg, J., Ek-Rylander, B. & Andersson, G. Tartrate-resistant purple acid phosphatase is synthesized as a latent proenzyme and activated by cysteine proteinases. *Biochem. J.* **343 Pt 1**, 63–9 (1999).
135. Oddie, G. W. *et al.* Structure , Function , and Regulation of Tartrate-Resistant Acid Phosphatase. *Bone* **27**, 575–584 (2000).

136. Fagerlund, K. M. *et al.* Effects of proteolysis and reduction on phosphatase and ROS-generating activity of human tartrate-resistant acid phosphatase. *Arch. Biochem. Biophys.* **449**, 1–7 (2006).
137. Wang, Y., Norgard, M., Andersson, G. & Norgård, M. N-glycosylation influences the latency and catalytic properties of mammalian purple acid phosphatase. *Arch. Biochem. Biophys.* **435**, 147–56 (2005).
138. Lång, P. *et al.* Monomeric tartrate resistant acid phosphatase induces insulin sensitive obesity. *PLoS One* **3**, e1713 (2008).
139. Hayman, a R., Bune, a J. & Cox, T. M. Widespread expression of tartrate-resistant acid phosphatase (Acp 5) in the mouse embryo. *J. Anat.* **196** (Pt 3), 433–41 (2000).
140. Klobusická, M. & Kusenda, J. Cytochemical study of activated peritoneal macrophages in normal and tumor-bearing rats. *Neoplasma* **39**, 291–7 (1992).
141. Hayman, a R., Macary, P., Lehner, P. J. & Cox, T. M. Tartrate-resistant acid phosphatase (Acp 5): identification in diverse human tissues and dendritic cells. *J. Histochem. Cytochem.* **49**, 675–684 (2001).
142. Bune, A. J., Hayman, A. R., Evans, M. J. & Cox, T. M. Mice lacking tartrate-resistant acid phosphatase (Acp 5) have disordered macrophage inflammatory responses and reduced clearance of the pathogen, *Staphylococcus aureus*. *Immunology* **102**, 103–113 (2001).
143. Esfandiari, E. *et al.* TRACP Influences Th1 pathways by affecting dendritic cell function. *J. Bone Miner. Res.* **21**, 1367–76 (2006).
144. Bevilacqua, M. A. *et al.* Regulation and expression of type V (tartrate-resistant) acid phosphatase in human mononuclear phagocytes. *Mol. Biol. Med.* **8**, 135–40 (1991).
145. Muhonen, P. *et al.* Sequence and TLR9 independent increase of TRACP expression by antisense DNA and siRNA molecules. *Biochem. Biophys. Res. Commun.* **359**, 889–95 (2007).
146. Lång, P., Lange, S., Delbro, D. & Andersson, G. Induction and cellular expression of tartrate resistant acid phosphatase during dextran sodium sulphate induced colitis in rats. *Histochem. Cell Biol.* **132**, 599–612 (2009).
147. Zenger, S. *et al.* Proteolytic processing and polarized secretion of tartrate-resistant acid phosphatase is altered in a subpopulation of metaphyseal osteoclasts in cathepsin K-deficient mice. *Bone* **41**, 820–32 (2007).
148. Lång, P. & Andersson, G. Differential expression of monomeric and proteolytically processed forms of tartrate-resistant acid phosphatase in rat tissues. *Cell. Mol. Life Sci.* **62**, 905–18 (2005).
149. Yamamoto, T. & Nagai, H. Ultrastructural localization of tartrate-resistant acid phosphatase activity in rat osteoblasts. *J. Electron Microsc. (Tokyo)*. **47**, 659–63 (1998).

150. Bianco, P., Ballanti, P. & Bonucci, E. Tartrate-resistant acid phosphatase activity in rat osteoblasts and osteocytes. *Calcif. Tissue Int.* **43**, 167–71 (1988).
151. Perez-Amodio, S. *et al.* Endogenous expression and endocytosis of tartrate-resistant acid phosphatase (TRACP) by osteoblast-like cells. *Bone* **36**, 1065–77 (2005).
152. Sheu, T.-J. J. *et al.* A phage display technique identifies a novel regulator of cell differentiation. *J. Biol. Chem.* **278**, 438–43 (2003).
153. Perez-Amodio, S., Jansen, D. C., Tigchelaar-Gutter, W., Beertsen, W. & Everts, V. Endocytosis of tartrate-resistant acid phosphatase by osteoblast-like cells is followed by inactivation of the enzyme. *Calcif. Tissue Int.* **78**, 248–54 (2006).
154. Solberg, L. B. *et al.* Increased tartrate-resistant Acid phosphatase expression in osteoblasts and osteocytes in experimental osteoporosis in rats. *Calcif. Tissue Int.* **94**, 510–21 (2014).
155. Gradin, P., Hollberg, K., Cassady, a I., Lång, P. & Andersson, G. Transgenic Overexpression of Tartrate-Resistant Acid Phosphatase Is Associated with Induction of Osteoblast Gene Expression and Increased Cortical Bone Mineral Content and Density. *Cells. Tissues. Organs* **196**, 68-81 (2012).
156. Solberg, L. B., Stang, E., Brorson, S.-H., Andersson, G. & Reinholt, F. P. Tartrate-resistant acid phosphatase (TRAP) co-localizes with receptor activator of NF- κ B ligand (RANKL) and osteoprotegerin (OPG) in lysosomal-associated membrane protein 1 (LAMP1)-positive vesicles in rat osteoblasts and osteocytes. *Histochem. Cell Biol.* **143**, 195–207 (2015).
157. Hollberg, K., Hultenby, K., Hayman, A., Cox, T. & Andersson, G. Osteoclasts from mice deficient in tartrate-resistant acid phosphatase have altered ruffled borders and disturbed intracellular vesicular transport. *Exp Cell Res* **279**, 227–238 (2002).
158. Lång, P. *et al.* Monomeric tartrate resistant Acid phosphatase induces insulin sensitive obesity. *PLoS One* **3**, e1713 (2008).
159. Lang, P., Schultzberg, M. & Andersson, G. Expression and Distribution of Tartrate-resistant Purple Acid Phosphatase in the Rat Nervous System. *J. Histochem. Cytochem.* **49**, 379–396 (2001).
160. Hayman, A. R., Bune, A. J., Bradley, J. R., Rashbass, J. & Cox, T. M. Osteoclastic Tartrate-resistant Acid Phosphatase (Acp 5): Its Localization to Dendritic Cells and Diverse Murine Tissues. *J Histochem Cytochem.* **48**, 219–227 (2015).
161. Ariyoshi, W. *et al.* Heparin inhibits osteoclastic differentiation and function. *J. Cell. Biochem.* **103**, 1707–17 (2008).
162. Igarashi Y, Lee MY, M. S. Heparin column analysis of serum type 5 tartrate-resistant acid phosphatase isoforms. *J Chromatogr B Biomed Sci Appl* **757** (2), 269–276 (2001).
163. Nakanishi, M. *et al.* Development of a kinetic assay for band 5b tartrate-resistant acid phosphatase activity in serum. *Clin. Chem.* **46**, 469–73 (2000).

164. Sheu, T.-J., Schwarz, E. M., O'Keefe, R. J., Rosier, R. N. & Puzas, J. E. Use of a phage display technique to identify potential osteoblast binding sites within osteoclast lacunae. *J. Bone Miner. Res.* **17**, 915–22 (2002).
165. Puzas, J. E., Sheu, T. J., Schwarz, E. M., O'Keefe, R. J. & Rosier, R. N. An osteoblast phage display library identifies TRIP to have high affinity for TRAP. *J Bone Min. Res* **16**, S164 (2001).
166. Metz-Estrella, D., Jonason, J. H., Sheu, T.-J., Mroczek-Johnston, R. M. & Puzas, J. E. TRIP-1: a regulator of osteoblast function. *J. Bone Miner. Res.* **27**, 1576–84 (2012).
167. Brehme, C. S., Roman, S., Shaffer, J. & Wolfert, R. Tartrate-resistant acid phosphatase forms complexes with alpha2-macroglobulin in serum. *J Bone Min. Res* **14**, 311–318 (1999).
168. Takahashi, K. *et al.* Electrophoretic study of tartrate-resistant acid phosphatase isoforms in endstage renal disease and rheumatoid arthritis. *Clin. Chim. Acta.* **301**, 147–58 (2000).
169. Janckila, A. J., Slone, S. P., Lear, S. C., Martin, A. & Yam, L. T. Tartrate-resistant acid phosphatase as an immunohistochemical marker for inflammatory macrophages. *Am. J. Clin. Pathol.* **127**, 556–66 (2007).
170. Janckila, A. J. *et al.* Serum tartrate-resistant acid phosphatase isoform 5a (TRACP5a) as a potential risk marker in cardiovascular disease. *Clin. Chim. Acta.* **412**, 963–9 (2011).
171. Janckila, A. J. *et al.* Serum tartrate-resistant acid phosphatase isoforms in rheumatoid arthritis. *Clin Chim Acta* **320**, 49–58 (2002).
172. Janckila, A. J., Neustadt, D. H. & Yam, L. T. Significance of Serum TRACP in Rheumatoid Arthritis. *J. Bone Miner. Res.* **23**, (2008).
173. Wu, Y.-Y., Janckila, A. J., Slone, S. P., Perng, W.-C. & Chao, T.-Y. Tartrate-resistant acid phosphatase 5a in sarcoidosis: further evidence for a novel macrophage biomarker in chronic inflammation. *J. Formos. Med. Assoc.* **113**, 364–70 (2014).
174. Briggs, T. a *et al.* Tartrate-resistant acid phosphatase deficiency causes a bone dysplasia with autoimmunity and a type I interferon expression signature. *Nat. Genet.* **43**, 127–131 (2011).
175. Ek-Rylander, B., Flores, M., Wendel, M., Heinegård, D. & Andersson, G. Dephosphorylation of osteopontin and bone sialoprotein by osteoclastic tartrate-resistant acid phosphatase. Modulation of osteoclast adhesion in vitro. *J. Biol. Chem.* **269**, 14853–6 (1994).
176. Andersson, G. *et al.* TRACP as an osteopontin phosphatase. *J. Bone Miner. Res.* **18**, 1912–5 (2003).

177. Lausch, E. *et al.* Genetic deficiency of tartrate-resistant acid phosphatase associated with skeletal dysplasia, cerebral calcifications and autoimmunity. *Nat. Genet.* **43**, 132–137 (2011).
178. Xia, L. *et al.* ACP5, a direct transcriptional target of FoxM1, promotes tumor metastasis and indicates poor prognosis in hepatocellular carcinoma. *Oncogene* **33**, 1395–406 (2014).
179. Adams, L., Warburton, M. & Hayman, A. Human breast cancer cell lines and tissues express tartrate-resistant acid phosphatase (TRAP). *Cell Biol. Int.* **31**, 191–195 (2007).
180. Honig, A. *et al.* Increased tartrate-resistant acid phosphatase (TRAP) expression in malignant breast, ovarian and melanoma tissue: an investigational study. *BMC Cancer* **6**, 199 (2006).
181. Kawamura, M. *et al.* Clinical significance of tartrate-resistant acid phosphatase type-5 expression in human gastric cancer. *Anticancer Res.* **34**, 3425–9 (2014).
182. How, J., Brown, J. R., Saylor, S. & Rimm, D. L. Macrophage expression of tartrate-resistant acid phosphatase as a prognostic indicator in colon cancer. *Histochem. Cell Biol.* **142**, 195–204 (2014).
183. Zenger, S. *et al.* Differential expression of tartrate-resistant acid phosphatase isoforms 5a and 5b by tumor and stromal cells in human metastatic bone disease. *Clin. Exp. Metastasis* **28**, 65–73 (2011).
184. Zenger, S., Ek-Rylander, B. & Andersson, G. Biogenesis of tartrate-resistant acid phosphatase isoforms 5a and 5b in stably transfected MDA-MB-231 breast cancer epithelial cells. *Biochim. Biophys. Acta* **1803**, 598–607 (2010).
185. Scott, K. L. *et al.* Proinvasion metastasis drivers in early-stage melanoma are oncogenes. *Cancer Cell* **20**, 92–103 (2011).
186. Simpson, K. J. *et al.* Identification of genes that regulate epithelial cell migration using an siRNA screening approach. *Nat. Cell Biol.* **10**, 1027–38 (2008).
187. Souza Malaspina, T. S. *et al.* Tartrate-resistant acid phosphatase activity and glutathione levels are modulated during hFOB 1.19 osteoblastic differentiation. *J. Mol. Histol.* **39**, 627–34 (2008).
188. Bazer, F. W., Worthington-White, D., Fliss, M. F. V & Gross, S. Uteroferrin: a progesterone-induced hematopoietic growth factor of uterine origin. *Exp. Hematol.* **19**, 910–915 (1991).
189. Ying, W., Wang, H., Bazer, F. W. & Zhou, B. Pregnancy-secreted Acid phosphatase, uteroferrin, enhances fetal erythropoiesis. *Endocrinology* **155**, 4521–30 (2014).
190. Ek-Rylander, B. & Andersson, G. Osteoclast migration on phosphorylated osteopontin is regulated by endogenous tartrate-resistant acid phosphatase. *Exp. Cell Res.* **316**, 443–51 (2010).

191. Al-Shami, R. *et al.* Phosphorylated osteopontin promotes migration of human choriocarcinoma cells via a p70 S6 kinase-dependent pathway. *J. Cell. Biochem.* **94**, 1218–33 (2005).
192. Lang, P. *et al.* Expression and secretion of the novel adipokine tartrate-resistant acid phosphatase from adipose tissues of obese and lean women. *Int. J. Obes. (Lond).* **35**, 1502–1510 (2011).
193. Shih, K.-C. *et al.* Effects of exercise on insulin sensitivity, inflammatory cytokines, and serum tartrate-resistant acid phosphatase 5a in obese Chinese male adolescents. *Metabolism.* **59**, 144–51 (2010).
194. Janckila, A. J. *et al.* Properties and expression of human tartrate-resistant acid phosphatase isoform 5a by monocyte-derived cells. *J Leukoc Biol* **77**, 209–218 (2005).
195. Chao, T. Y. *et al.* Development of immunoassays for serum tartrate-resistant acid phosphatase isoform 5a. *Clin Chim Acta* **359**, 132–140 (2005).
196. Kimura, N., Toyoshima, T., Kojima, T. & Shimane, M. Entactin-2: a new member of basement membrane protein with high homology to entactin/nidogen. *Exp. Cell Res.* **241**, 36–45 (1998).
197. Kohfeldt, E., Sasaki, T., Göhring, W. & Timpl, R. Nidogen-2: a new basement membrane protein with diverse binding properties. *J. Mol. Biol.* **282**, 99–109 (1998).
198. Rahman, S. *et al.* Novel hepatocyte growth factor (HGF) binding domains on fibronectin and vitronectin coordinate a distinct and amplified Met-integrin induced signalling pathway in endothelial cells. *BMC Cell Biol.* **6**, 8 (2005).
199. Walker PD, Kaushal GP, Shah SV. Meprin A , the major matrix degrading enzyme in renal tubules , produces a novel nidogen fragment in vitro and in vivo. *Kidney Int.* **53**, 1673–1680 (1998).
200. Chen, R., Miettinen, P., Marouka, E., Choy, L. & Derynck, R. A WD-domain protein that is associated with and phosphorylated by the type II TGF beta receptor. *Nature* **377**, 548–552 (1995).
201. Sieber, J. J. *et al.* Anatomy and dynamics of a supramolecular membrane protein cluster. *Science* **317**, 1072–6 (2007).
202. Yayon, a, Klagsbrun, M., Esko, J. D., Leder, P. & Ornitz, D. M. Cell surface, heparin-like molecules are required for binding of basic fibroblast growth factor to its high affinity receptor. *Cell* **64**, 841–8 (1991).
203. Rapraeger, a C., Krufka, a & Olwin, B. B. Requirement of heparan sulfate for bFGF-mediated fibroblast growth and myoblast differentiation. *Science* **252**, 1705–8 (1991).
204. Hsu, Y. R. *et al.* Heparin is essential for a single keratinocyte growth factor molecule to bind and form a complex with two molecules of the extracellular domain of its receptor. *Biochemistry* **38**, 2523–34 (1999).

205. Schaefer, L. & Schaefer, R. M. Proteoglycans: from structural compounds to signaling molecules. *Cell Tissue Res.* **339**, 237–46 (2010).
206. Häcker, U., Nybakken, K. & Perrimon, N. Heparan sulphate proteoglycans: the sweet side of development. *Nat. Rev. Mol. Cell Biol.* **6**, 530–541 (2005).
207. Lajoie, P. & Nabi, I. R. *Lipid rafts, caveolae, and their endocytosis*. *Int Rev Cell Mol Biol.* **282**, 135–63 (2010).
208. Gustavsson, J. *et al.* Localization of the insulin receptor in caveolae of adipocyte plasma membrane. *FASEB J.* **13**, 1961–71 (1999).
209. Ost, A., Ortegren, U., Gustavsson, J., Nystrom, F. H. & Strålfors, P. Triacylglycerol is synthesized in a specific subclass of caveolae in primary adipocytes. *J. Biol. Chem.* **280**, 5–8 (2005).
210. Chen, S. *et al.* Wnt-1 Signaling Inhibits Apoptosis by Activating β -Catenin / T Cell Factor – mediated Transcription. *J Cell Biol.* **152**, 87–96 (2001).
211. Longo, K. a *et al.* Wnt signaling protects 3T3-L1 preadipocytes from apoptosis through induction of insulin-like growth factors. *J. Biol. Chem.* **277**, 38239–44 (2002).
212. Calò, V. *et al.* STAT proteins: from normal control of cellular events to tumorigenesis. *J. Cell. Physiol.* **197**, 157–68 (2003).
213. Harp, J. B. Activation of Signal Transducer and Activator of Transcription-3 during Proliferative Phases of 3T3-L1 Adipogenesis . *Endocrinology* **141**, 2370–2376 (2000).
214. Venkatakrishnan, C. D. *et al.* HSP27 regulates p53 transcriptional activity in doxorubicin-treated fibroblasts and cardiac H9c2 cells: p21 upregulation and G2/M phase cell cycle arrest. *Am. J. Physiol. Heart Circ. Physiol.* **294**, H1736–44 (2008).
215. Rousseau, S. *et al.* CXCL12 and C5a trigger cell migration via a PAK1/2-p38alpha MAPK-MAPKAP-K2-HSP27 pathway. *Cell. Signal.* **18**, 1897–905 (2006).
216. Page, K., Li, J. & Hershenson, M. B. p38 MAP kinase negatively regulates cyclin D1 expression in airway smooth muscle cells. *Am J Physiol Lung Cell Mol Physiol.* **280**, 955–964 (2001).
217. Sun, P. *et al.* PRAK is essential for ras-induced senescence and tumor suppression. *Cell* **128**, 295–308 (2007).
218. Serrano, M., Lin, A. W., Mccurrach, M. E., Beach, D. & Lowe, S. W. Oncogenic ras Provokes Premature Cell Senescence Associated with Accumulation of p53 and p16 INK4a. **88**, 593–602 (1997).
219. Roger, L., Gadea, G. & Roux, P. Control of cell migration: a tumour suppressor function for p53? *Biol. Cell* **98**, 141–52 (2006).

220. Kobayashi, M., Nishita, M., Mishima, T., Ohashi, K. & Mizuno, K. MAPKAPK-2-mediated LIM-kinase activation is critical for VEGF-induced actin remodeling and cell migration. *EMBO J.* **25**, 713–26 (2006).
221. Del Barco Barrantes, I. & Nebreda, A. R. Roles of p38 MAPKs in invasion and metastasis. *Biochem. Soc. Trans.* **40**, 79–84 (2012).
222. Maged, F., Gurgis, S., Ziaziaris, W. & Munoz, L. Mitogen-Activated Protein Kinase – Activated Protein Kinase 2 in Neuroinflammation , Heat Shock Protein 27 Phosphorylation , and Cell Cycle : Role and Targeting. *Mol Pharmacol.* **85**, 345–356 (2014).
223. Corcoran, A. & Del Maestro, R. F. Testing the ‘Go or Grow’ Hypothesis in Human Medulloblastoma Cell Lines in Two and Three Dimensions. *Neurosurgery* **53**, 174–185 (2003).
224. Garay, T. *et al.* Cell migration or cytokinesis and proliferation?--revisiting the ‘go or grow’ hypothesis in cancer cells in vitro. *Exp. Cell Res.* **319**, 3094–103 (2013).
225. Ek-Rylander, B. & Andersson, G. Osteoclast migration on phosphorylated osteopontin is regulated by endogenous tartrate-resistant acid phosphatase. *Exp Cell Res* **316**, 443–451 (2010).
226. Magun, R., Boone, D. L., Tsang, B. K. & Sorisky, A. The effect of adipocyte differentiation on the capacity of 3T3-L1 cells to undergo apoptosis in response to growth factor deprivation. *Int J Obes Relat Metab Disord* **22**, 567-71 (1998).
227. Sakane, H., Yamamoto, H., Matsumoto, S., Sato, A. & Kikuchi, A. Localization of glypican-4 in different membrane microdomains is involved in the regulation of Wnt signaling. *J. Cell Sci.* **125**, 449–60 (2012).